
**Soil quality — Determination of soil
microbial biomass —**

**Part 2:
Fumigation-extraction method**

*Qualité du sol — Détermination de la biomasse microbienne du sol —
Partie 2: Méthode d'extraction par fumigation*

STANDARDSISO.COM : Click to view the full PDF of ISO 14240-2:1997



Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

International Standard ISO 14240-2 was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

ISO 14240 consists of the following parts, under the general title *Soil quality — Determination of soil microbial biomass*:

- *Part 1: Substrate-induced respiration method*
- *Part 2: Fumigation-extraction method*

STANDARDSISO.COM : Click to view the full PDF of ISO 14240-2:1997

© ISO 1997

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the publisher.

International Organization for Standardization
Case postale 56 • CH-1211 Genève 20 • Switzerland
Internet central@iso.ch
X.400 c=ch; a=400net; p=iso; o=isocs; s=central

Printed in Switzerland

Introduction

Soil consists of both living and nonliving components which exist in a complex and heterogeneous environment. Soil microflora is responsible for the degradation of organic matter, stability of aggregates and most nutrient cycling which occurs in soils. The purpose of determining the microbial biomass of soils is to allow assessment of the continued maintenance of soil fertility, the potential ability to degrade added organic materials, and the effects of added materials on the natural microbial population.

STANDARDSISO.COM : Click to view the full PDF of ISO 14240-2:1997

This page intentionally left blank

STANDARDSISO.COM : Click to view the full PDF of ISO 14240-2:1997

Soil quality — Determination of soil microbial biomass —

Part 2: Fumigation-extraction method

1 Scope

This part of ISO 14240 specifies a method for the estimation of microbial biomass of soils by measurement of total extractable organic biomass material mainly from freshly killed microorganisms. The method is also applicable to the estimation of microbial nitrogen and microbial ninhydrin-reactive nitrogen in soil, but this part of ISO 14240 describes only the measurement of extractable organic carbon. The fumigation-extraction (FE) method is applicable to aerobic and anaerobic (water-logged, paddy) soils over the whole range of soil pH. Biomass can be also measured in soils containing actively decomposing substrates and soils supersaturated with potassium sulfate solution.

NOTE — Chloroform fumigation also affects soil fauna. The contribution of carbon from these organisms is generally small (< 5 %) and can usually be neglected.

2 Normative references

The following standards contain provisions which, through reference in this text, constitute provisions of this part of ISO 14240. At the time of publication, the editions indicated were valid. All standards are subject to revision, and parties to agreements based on this part of ISO 14240 are encouraged to investigate the possibility of applying the most recent editions of the standards indicated below. Members of IEC and ISO maintain registers of currently valid International Standards.

ISO 10381-6:1993, *Soil quality — Sampling — Part 6: Guidance on the collection, handling and storage of soil for the assessment of aerobic microbial processes in the laboratory.*

ISO 10694:1995, *Soil quality — Determination of organic and total carbon after dry combustion (elementary analysis).*

ISO 11465:1993, *Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method.*

3 Definition

For the purposes of this part of ISO 14240, the following definition applies.

3.1 soil microbial biomass

mass of intact microbial cells in a given soil

NOTE — This parameter can be estimated from the measurement of the carbon or nitrogen content of these cells or by the measurement of their ability to mineralize an added carbon source. Dead cells and cell fragments may be detected when carbon or nitrogen analysis is used but only intact cells will be detected when respiration is measured.

4 Principle

Through fumigation of the soil sample, intact microbial cells are lysed and the microbial organic matter released. Nonliving soil organic matter is not seriously affected by such fumigation. Soil samples are fumigated with chloroform for 24 h. The organic carbon extracted using 0,5 mol/l potassium sulfate is determined in fumigated and unfumigated samples, and the difference in extracted organic carbon is used to determine microbial biomass carbon. The method is usually termed fumigation-extraction (FE).

5 Materials and reagents

5.1 Soils

Guidance for the collection, handling and storage of soil (ISO 10381-6) shall be followed, as far as applicable. Sieve the soil samples at approximately 40 % water-holding capacity (WHC) if homogeneous samples are required.

Determine the soil WHC in accordance with annex A.

The water content of samples shall be higher than 30 % WHC to ensure uniform chloroform distribution and effective fumigation. Take care to avoid smearing and compaction of wet soil in this method. Samples of waterlogged soils do not have to be dried prior to analysis.

5.2 Reagents

Reagents of recognized analytical grade shall be used, including:

5.2.1 Silicone grease (medium viscosity).

5.2.2 Ethanol-free chloroform.

WARNING — In the presence of light, ethanol-free chloroform degrades rapidly to form phosgene gas (COCl_2) which is odourless and highly toxic.

5.2.3 Potassium sulfate solution, $c(\text{K}_2\text{SO}_4) = 0,5 \text{ mol/l}$ ($\rho = 87,135 \text{ g/l}$).

5.2.4 Soda lime.

6 Apparatus

6.1 Room, or incubator, capable of being maintained at $(25 \pm 2) \text{ }^\circ\text{C}$.

6.2 Implosion-protected desiccator.

6.3 Filter paper¹⁾.

6.4 Glass beakers.

6.5 Petri dishes.

6.6 Polyethylene bottles, of capacity 250 ml.

1) Whatman No. 42, Schleicher & Schüll 595 1/2, Machery & Nagel 261 G 1/4, are examples of suitable products available commercially. This information is given for the convenience of users of this part of ISO 14240 and does not constitute an endorsement by ISO of these products.

6.7 Vacuum line (water-pump or electric pump).

6.8 Horizontal or overhead shaker.

6.9 Freezer, operation at (–15 to –20) °C.

6.10 Antibumping granules.

7 Fumigation and extraction

7.1 Fumigation

Line the desiccator (6.2) with moist filter paper (6.3) for fumigation of the soil samples.

Weigh at least three samples of moist soil (5.1), each containing a mass equivalent to 25 to 50 g of oven-dry soil, into glass beakers (6.4) or Petri dishes (6.5) and place them in the desiccator with a beaker containing 25 ml of ethanol-free chloroform (5.2.2) and a few antibumping granules (6.10), and a beaker with soda lime (5.2.4). Evacuate the desiccator until the chloroform has boiled vigorously for approximately 2 min. Close the vacuum tap on the desiccator and incubate in the dark (6.1) at (25 ± 2) °C for 22 h to 24 h.

If there is insufficient soil available, use smaller samples provided the ratio of soil mass to volume of extractant is kept the same (1:4). In soils containing more than 20 % organic material (as determined according to ISO 10694), increase the ratio soil:extract above 1:4 (to a maximum of 1:30 for soils containing 95 % organic matter, e.g. L-layer) in order to obtain sufficient extracted matter. Record the mass of soil used.

After fumigation is completed, remove the beaker containing the chloroform and the filter paper from the desiccators. Remove the chloroform vapour from the soil by repeated evacuation (6 times for 2 min each) in the desiccator. The samples are ready for extraction.

Place three unfumigated, moist control samples (50 g dry mass) into polyethylene bottles (6.6) and immediately extract with 200 ml of the potassium sulfate (5.2.3), as indicated in 7.2.

7.2 Extraction

To extract organic carbon, transfer the soil quantitatively to polyethylene bottles (6.6), add 200 ml of potassium sulfate (5.2.3), shake the bottles on a horizontal shaker (6.8) at 200 r/min for 30 min or an overhead shaker at 60 r/min for 45 min and filter the extracts through a folded filter paper (6.3). Extract the unfumigated controls and filter the extracts in the same way.

If not analyzed at once, store the extracts of fumigated and unfumigated soil samples in the freezer (6.9) at between –15 °C and –20 °C. Homogenize frozen extracts before use, after thawing at room temperature.

NOTES

1 A white precipitate occurs during storage of the potassium sulfate soil extracts (especially if the samples are frozen) because they are usually supersaturated with calcium sulfate (CaSO_4). It is unnecessary to dissolve this excess CaSO_4 because it does not interfere with any of the analytical procedures described in this method.

2 Cell membranes of young, living roots are also affected by chloroform fumigation. In soils containing large amounts of living roots, the pre-extraction procedure given in annex B should be used.

8 Determination of carbon in the extracts

The carbon content of microorganisms in a soil sample is determined analytically and can be used to make comparisons between different soil samples. Where a figure for actual microbial biomass is required, then such analyses are multiplied by a conversion factor derived from experiments correlating a known cell mass to carbon analysis after fumigation-extraction. All conversion factors used are related to this initial factor.

Determine the carbon content in the extracts using either dichromate oxidation (8.1) or instrumental analysis (8.2).

8.1 Microbial biomass carbon by dichromate oxidation

8.1.1 Principle

In the presence of strong acid, organic matter is oxidized and Cr(VI) reduced to Cr(III). The amount of dichromate left is back-titrated.

8.1.2 Additional reagents

8.1.2.1 Potassium dichromate, $c(\text{K}_2\text{Cr}_2\text{O}_7) = 0,0667 \text{ mol/l}$ (19,6125 g dry potassium dichromate/litre water).

WARNING — Because of the hazardous nature of potassium dichromate, great care should be taken in its use and eventual disposal.

8.1.2.2 Phosphoric acid, (H_3PO_4) , $\rho = 1,71 \text{ g/ml}$.

8.1.2.3 Sulfuric acid, (H_2SO_4) , $\rho = 1,84 \text{ g/ml}$.

8.1.2.4 Iron(II) ammonium sulfate, titration solution, $c[(\text{NH}_4)_2\text{Fe}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}] = 0,040 \text{ mol/l}$.

Iron(II) ammonium sulfate (15,69 g) is dissolved in distilled water, acidified with 20 ml of the sulfuric acid (8.1.2.3) and made up to 1 000 ml with distilled water.

8.1.2.5 1,10-phenanthroline sulfate complex solution, 0,025 mol/l.

8.1.2.6 Acid mixture: two volumes sulfuric acid (8.1.2.3) mixed with one volume phosphoric acid (8.1.2.2).

8.1.3 Additional apparatus

8.1.3.1 Liebig condenser (water-cooled).

8.1.3.2 250 ml round-bottom flask.

8.1.3.3 Burette, volume 10 ml, graduated in 0,05 ml divisions.

8.1.3.4 Pipette, volume 2 ml.

8.1.4 Procedure

Add 2 ml of the potassium dichromate solution (8.1.2.1) (P_D) using the pipette (8.1.3.4) and 15 ml of the acid mixture (8.1.2.6) to 8 ml of filtered extract (7.1) (P_S) in a 250 ml round-bottom flask (8.1.3.2). Gently reflux (8.1.3.1) the whole mixture for 30 min, allow to cool and dilute with 20 ml to 25 ml water, added through the condenser to rinse it.

Digest duplicate blanks containing 8 ml of the potassium sulfate solution (5.2.3) in the same way.

NOTE — These digested blanks are also known as "refluxed blanks".

Determine the excess dichromate by back-titration (8.1.3.3) with the iron(II) ammonium sulfate titration solution (8.1.2.4), using a few drops of the 1,10-phenanthroline-iron(II) sulfate complex solution (8.1.2.5) as an indicator.

8.1.5 Calculation of results

Calculate the extractable organic carbon using equations (1) and (2).

$$C (\mu\text{g/ml}) = [(V_H - V_S) / V_C] \times MP_D \times E \times 1000 / P_S \quad \dots (1)$$

where

V_s is the volume titrant (8.1.2.4) consumed by the sample, in millilitres;

V_H is the volume titrant (8.1.2.4) consumed by the refluxed blank, in millilitres;

V_C is the volume titrant (8.1.2.4) consumed by the unrefluxed blank, in millilitres;

M is $c(K_2Cr_2O_7)$, in moles per litre;

P_D is the added volume of $K_2Cr_2O_7$ solution, in millilitres;

P_s is the added volume of sample, in millilitres;

E is 3{conversion of organic C[C] to $CO_2[C(+IV)]$ }.

$$C (\mu\text{g} / \text{g dry soil}) = C (\mu\text{g} / \text{ml}) \times (P_K/D_w + S_w) \quad \dots (2)$$

where

P_K is the mass of extractant, in grams;

D_w is the dry mass of sample (determined according to ISO 11465), in grams;

S_w is the soil water (grams of water/grams of dry soil) (determined according to ISO 11465).

Calculate the biomass carbon, B_c , using equation (3).

$$B_c = E_c / k_{EC} \quad \dots (3)$$

where

E_c = (mass of organic C extracted from fumigated soils) – (mass of organic C extracted from unfumigated soils);

$k_{EC} = 0,38$.

NOTE — The factor k_{EC} was calculated by relating the results of the fumigation-incubation method and fumigation-extraction method (12 soils).

8.2 Microbial biomass carbon by spectrometric carbon analysis

8.2.1 Principle

In the presence of potassium persulfate ($K_2S_2O_8$), extractable soil organic carbon is oxidized to carbon dioxide, and measured using infrared (IR) or ultraviolet (UV) spectrometric detection.

8.2.2 Additional reagents

8.2.2.1 Potassium persulfate ($K_2S_2O_8$).

8.2.2.2 Phosphoric acid (see 8.1.2.2).

8.2.2.3 Sodium polyphosphate [$(NaPO_3)_n$], extra pure.

8.2.2.4 Potassium persulfate reagent

Dissolve 20 g of potassium persulfate (8.2.2.1) in 900 ml of distilled water, acidify the solution to pH 2 with phosphoric acid (8.2.2.2) and make up to a final volume of 1 000 ml with distilled water.

8.2.2.5 Sodium polyphosphate reagent

Dissolve 50 g of sodium polyphosphate (8.2.2.3) in 900 ml of distilled water, acidify the solution to pH 2 with phosphoric acid (8.2.2.2) and make up to a final volume of 1 000 ml with distilled water.

8.2.3 Additional apparatus

8.2.3.1 Automatic carbon analyzer with IR detection¹⁾ or continuous-flow system with colorimetric detection²⁾.

For the purposes of this part of ISO 14240, the determination of microbial biomass carbon is based on UV-activated persulfate oxidation of organic carbon.

8.2.4 Procedure

For the automated UV-persulfate oxidation method, mix 5 ml of the potassium sulfate soil extract (7.1) with 5 ml of sodium polyphosphate reagent (8.2.2.5). Any precipitate of CaSO₄ in the soil extracts is dissolved by this procedure. The potassium persulfate reagent (8.2.2.4) is automatically fed into the UV oxidation chamber, where the oxidation to CO₂ is activated by UV light. The resulting CO₂ is measured by IR absorption or UV spectrophotometric detection.

8.2.5 Calculation of results

Calculate the extractable organic carbon (*C*) using equation (4).

$$C (\mu\text{g} / \text{g dry soil}) = [(V \times D_v) - (B \times D_b)] \times (P_k / D_w + S_w) \quad \dots (4)$$

where

V is the *C* (μg / ml) of sample;

B is the *C* (μg / ml) of blank;

D_v is the dilution of sample with sodium phosphate, in millilitres;

D_b is the dilution of blank with sodium phosphate, in millilitres;

P_k see equation (2);

D_w see equation (2);

S_w see equation (2).

Calculate the biomass *B_c* using equation (5).

$$B_c = E_c / k_{ec} \quad \dots (5)$$

where

E_c = (organic C extracted from fumigated soils) – (organic C extracted from unfumigated soils);

k_{ec} = 0,45.

¹⁾ Dohrman DC 80, Maihak Tocor 4, are examples of suitable equipment available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of these products.

²⁾ Skalar, Perstorp Alpkem, are examples of suitable equipment available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of these products.

NOTES

- 1 The factor k_{ec} was calculated by relating the results of the fumigation-incubation method to those of fumigation-extraction method (23 soils).
- 2 It is possible to use the fumigation-extraction method in conjunction with decomposition studies of ^{14}C -labelled organic substrates.

9 Precision

Details of an interlaboratory test on the precision of the method are summarized in annex C. The values derived from this interlaboratory test may not be applicable to concentration ranges and matrices other than those given.

STANDARDSISO.COM : Click to view the full PDF of ISO 14240-2:1997

Annex A (normative)

Measurement of water-holding capacity of soil

A.1 Scope

The method is applicable for obtaining a value for the water-holding capacity of a soil for applications in which the exact value is not critical.

A.2 Principle

A cylinder with perforated base is partially filled with soil, capped and immersed in water and drained. The quantity of water taken up by the soil is determined by weighing, drying to constant mass at 105 °C and reweighing.

A.3 Apparatus

A.3.1 Cylinder, of known volume, of approximate dimensions 50 mm to 150 mm in length and 50 mm to 100 mm in diameter, with a perforated base.

A.3.2 Water bath (room temperature).

A.3.3 Tray, with a drainage hole, containing wet fine quartz sand to a depth of 20 mm to 50 mm.

A.3.4 Oven, capable of maintaining a temperature of 105 °C ± 2 °C.

A.3.5 Balance, capable of weighing to an accuracy of ± 0,01 g.

A.4 Procedure

Cover the perforated base of the cylinder (A.3.1) with a filter paper and weigh the cylinder with the filter paper. Partially fill the cylinder with soil and cap the cylinder. Submerge the cylinder in a water bath for 2 h at room temperature with the water level for one hour. Remove the cylinder from the water and place the cylinder on a tray of sand (A.3.3) and allow to drain for 2 h to 24 h, depending on the soil type. Weigh the cylinder with soil, remove the soil and dry it to constant mass at 105 °C, and weigh the soil again.

A.5 Calculation

Calculate the water-holding capacity *WHC*, expressed as a percentage, as follows:

$$WHC = \frac{S - T - D}{D} \times 100$$

where

S is the mass of water-saturated soil + cylinder + filter paper, in grams;

T is the tare (mass of cylinder + filter paper), in grams;

D is the dry mass of soil, in grams.

A.6 Expression of results

Express the water-holding capacity of the soil (*WHC*) as a percentage of the dry mass of soil.

STANDARDSISO.COM : Click to view the full PDF of ISO 14240-2:1997

Annex B (informative)

Estimation of soil microbial biomass in soils containing large amounts of living roots — Pre-extraction procedure

B.1 Additional reagent

B.1.1 Potassium sulfate solution $c(\text{K}_2\text{SO}_4) = 0,05 \text{ mol/l}$; (8,714 g/l of extra-pure potassium sulfate).

B.2 Additional apparatus

B.2.1 Centrifuge.

B.3 Procedure

Place moist soil (25 g to 50 g dry mass) in a 250-ml glass bottle and pre-extract with 100 ml of the potassium sulfate solution (B.1.1) for 20 min by oscillation shaking at 200 r/min and pass through a sieve (for arable soils 2 mm mesh, for grassland soils 3 mm mesh). Carefully wash roots (and small stones) on the sieve until free of soil with an additional 75 ml of the potassium sulfate solution (B.1.1) and dry and weigh them. Centrifuge (B.2.1) the soil suspension in the original glass bottle for 15 min at ca. 500g. Then decant the supernatant. Add three drops of liquid chloroform (5.2.2) to soils to be fumigated. Proceed as in 7.1.

B.4 Comment

It is essential to use the procedure given above in the presence of living roots. In addition, it makes measurement of microbial biomass nitrogen easier by decreasing any background in inorganic nitrogen that the soil may contain. It also reduces the problems of measuring microbial biomass in dry soils and is therefore very suitable for measuring fluctuations in soil microbial biomass carbon and nitrogen during the year. No measurable microbial biomass is extracted from soils by the pre-extraction procedure.

Annex C (informative)

Results of an interlaboratory trial

The method was ring-tested in 11 laboratories in Germany. Two arable soils were used, namely a loess soil and a loam soil. The results given in table C.1 refer to organic carbon extracted from unfumigated and fumigated soils and E_c values. The E_c value is calculated from the difference between organic carbon extracted from fumigated soils minus organic carbon extracted from unfumigated soils. The biomass content of the soils is calculated by dividing E_c by the factor k_{EC} .

Table C.1 — Results of German interlaboratory ring test for the estimation of soil microbial biomass by the fumigation-extraction method

Parameter	Mean $\mu\text{g} / \text{g}$	CV ¹⁾ %	SE ²⁾
Loess soil			
C extracted from unfumigated samples	71	21	6.7
C extracted from fumigated samples	207	15	7.4
E_c ³⁾	136	23	9.2
Loam soil			
C extracted from unfumigated samples	85	18	5.4
C extracted from fumigated samples	265	11	7.1
E_c ³⁾	180	15	8.6
1) CV = coefficient of variation.			
2) SE = standard error of the mean ($m = 4$) of a single participant.			
3) Difference between results obtained for fumigated and unfumigated samples.			

Annex D (informative) Bibliography

- [1] ISO 10390:1994, *Soil quality — Determination of pH*.
- [2] ISO 11274:—¹⁾, *Soil quality — Determination of water-retention characteristics — Laboratory methods*.
- [3] BROOKES P.C., *et al.* Chloroform fumigation and the release of soil nitrogen: A rapid direct extraction method for measuring microbial biomass nitrogen in soil. *Soil Biol. Biochem.*, **17**, 1985, pp. 837-842.
- [4] HARDEN T., *et al.* Soil microbial biomass estimated by fumigation-extraction and substrate-induced respiration in two pesticide-treated soils. *Soil Biol. Biochem.*, **25**, 1993, pp. 679-683.
- [5] HARDEN T., *et al.* Mineralization of straw and formation of soil microbial biomass in a soil treated with Simazine and Dinoterb. *Soil Biol. Biochem.*, **25**, 1993, pp. 1273-1276.
- [6] INUBUSHI K., BROOKES P.C. and JENKINSON D.S. Soil microbial biomass C, N and ninhydrin-N in aerobic and anaerobic soils measured by the fumigation-extraction method. *Soil Biol. Biochem.*, **23**, 1991, pp. 737-741.
- [7] MUELLER T., JOERGENSEN R.G. and MEYER B. Estimation of soil microbial biomass C in the presence of living roots by fumigation-extraction, *Soil Biol. Biochem.*, **24**, 1992, pp. 179-181.
- [8] OCIO J.A. and BROOKES P.C. An evaluation of methods for measuring the microbial biomass in soils following recent additions of wheat straw and the characterization of the biomass that develops. *Soil Biol. Biochem.*, **22**, 1990, pp. 685-694.
- [9] SPARLING G.P., *et al.* Estimation of soil microbial C by a fumigation-extraction method: use on soils of high organic matter content, and a reassessment of the K_{ec} factor. *Soil Biol. Biochem.*, **22**, 1990, pp. 301-307.
- [10] WU J., *et al.* Measurement of soil microbial biomass C by fumigation-extraction: an automated procedure. *Soil Biol. Biochem.*, **22**, 1990, pp. 1167-1169.
- [11] VANCE E.D., BROOKES P.C. and JENKINSON D.S. An extraction method for measuring soil microbial C. *Soil Biol. Biochem.*, **19**, 1987, pp. 703-708.
- [12] WU J., BROOKES P.C. and JENKINSON D.S. Formation and destruction of microbial biomass during the decomposition of glucose and rye grass in soil. *Soil Biol. Biochem.*, **25**, 1993, pp. 1435-1441.

¹⁾ To be published.