
**Microbiology of the food chain —
Methods for the detection of
Anisakidae L3 larvae in fish and
fishery products —**

**Part 2:
Artificial digestion method**

*Microbiologie de la chaîne alimentaire — Méthodes de recherche des
larves L3 d'Anisakidae dans les poissons et produits de la pêche —*

Partie 2: Méthode de digestion artificielle

STANDARDSISO.COM : Click to view the full PDF of ISO 23036-2:2021



STANDARDSISO.COM : Click to view the full PDF of ISO 23036-2:2021



COPYRIGHT PROTECTED DOCUMENT

© ISO 2021

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

	Page
Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 Principle	2
4.1 General	2
4.2 Sample size	2
4.3 Sample preparation	2
4.4 Digestion of the sample	2
4.5 Filtration of the digest fluid	2
4.6 Verification of findings	2
5 Reagents	3
6 Equipment and consumables	3
7 Sampling	4
8 Procedure	4
8.1 Preparation of the sample	4
8.2 Preparation of the digest fluid	4
8.3 Digestion of the sample in the glass beaker	5
8.4 Filtration of the digest fluid	5
8.5 Microscopic examination	5
9 Expression of the results	5
10 Performance characteristics of the method	6
11 Test report	6
12 Quality assurance	6
Annex A (informative) Sample collection	7
Annex B (informative) Example of a laboratory worksheet for recording data when testing fish fillets with the artificial digestion method	8
Bibliography	9

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology*, in collaboration with the European Committee for Standardization (CEN) Technical Committee CEN/TC 275, *Food analysis — Horizontal methods*, in accordance with the Agreement on technical cooperation between ISO and CEN (Vienna Agreement).

A list of all parts in the ISO 23036 series can be found on the ISO website.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

Nematodes of the Anisakidae family have a complex life cycle involving a high number of hosts. Adult stages of Anisakidae reside in the stomach of marine mammals, where they are embedded in the mucosa. Unembryonated eggs produced by adult females are released with the faeces of marine mammals and become embryonated in seawater, where first-stage larvae (L1) develop in the eggs. The larvae moult to become free-swimming second-stage larvae (L2) and, if ingested by crustaceans, mature into third-stage larvae (L3). This stage is infective to fish and squid, and larvae are transferred between fishes through predation, maintaining the L3 stage. Some larvae migrate from the abdominal cavity into muscle tissues. Humans are incidental hosts and can be infected after ingesting raw or undercooked infected fish or cephalopods containing viable L3.

Nematodes of the family Anisakidae are the causative agents of human anisakidosis, a disease that is not only a public health hazard affecting humans, but also represents an economic problem in fishery and food safety (the term “anisakiasis”, designating the disease caused by members of the genus *Anisakis*, is also sometimes used). Worldwide, marine and wild anadromous fishes are intermediate hosts of Anisakidae, whereas marine mammals are the definitive hosts.

Visual inspection procedures for the detection of Anisakidae larvae in fish are employed to minimize the risk that contaminated fish will reach the consumer,^{[1][2]} thus preventing human anisakidosis.

The UV-press and the artificial digestion of the fish muscular tissue are the methods specifically designed to detect nematode larvae in fish and to evaluate the infestation level of a batch, and have been validated and tested in multicentre collaborative studies^[3] (see [Clause 10](#)).

STANDARDSISO.COM : Click to view the full PDF of ISO 23036-2:2021

Microbiology of the food chain — Methods for the detection of Anisakidae L3 larvae in fish and fishery products —

Part 2: Artificial digestion method

1 Scope

This document specifies a method for the detection of Anisakidae L3 larvae commonly found in marine and anadromous fishes. The method is applicable to fresh fish and/or frozen fish, as well as lightly processed fish products, such as marinated, salted or smoked. It is also suitable for visceral organs as a confirmatory method for a visual inspection scheme.

The artificial digestion method^{[4][5][6]} is applicable to quantifying parasitic infections by estimating the number of parasites in the fish musculature and, when applied to fresh fish or lightly processed fish products (never frozen before processing), determining the viability of Anisakidae L3, which can be present.

This method does not apply to determining the species or genotype of detected parasites. Final identification is made by morphological and/or molecular methods.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 7218, *Microbiology of food and animal feeding stuffs — General requirements and guidance for microbiological examinations*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <http://www.electropedia.org/>

3.1

Anisakidae L3 larvae

third-stage larvae (L3) belonging to the Anisakidae family, in particular to *Anisakis*, *Contracaecum* and *Pseudoterranova* genera

Note 1 to entry: For practical purposes, the genus *Hysterothylacium* belonging to Raphidascarididae family and already classified as Anisakidae, can be also included.

3.2 artificial digestion method

method to detect Anisakidae larvae in fish muscle tissue by an enzymatic (pepsin-HCl) digestion step to release larvae from muscle tissues followed by a filtration step and detection of larvae by microscopy

Note 1 to entry: If applied to fresh fish, it allows checking of the viability of larvae.

4 Principle

4.1 General

The artificial digestion method relies on the enzymatic degradation of muscle fibres in a fluid composed of pepsin and hydrochloric acid followed by filtration and washing steps.

The procedure allows a differentiation between dead and viable anisakid larvae if the temperature of the digestion solution does not exceed 37 °C (with the exception of *Hysterothylacium* sp. larvae, which are killed at 37 °C), and assuming the fish was never frozen.

There are no internal quality controls that can be used while performing the method.

NOTE Alternative methods can be used for analysis, provided their equivalence with the methods described in this document are demonstrated.

4.2 Sample size

For inspection of fishery products for public health purposes, the sample size shall be risk-based, as determined by the competent authority.

4.3 Sample preparation

To increase the surface area for enzymatic degradation, samples are gently eased apart taking care not to disrupt larvae by checking for them. Alternatively, a Smasher/Stomacher¹⁾, that facilitates the digestion but does not damage the nematode larvae, can be used.

A blending or grinding procedure should be avoided as this can damage or disrupt larvae.

4.4 Digestion of the sample

Viable Anisakidae larvae are resistant to the pepsin-HCl digest fluid and therefore can be recovered free from muscle tissues.

To facilitate an efficient and rapid digestion, a maximum ratio of 1:20, meat to digest fluid, and a temperature of 37 °C ± 2 °C shall be maintained throughout the process. The time required for digestion shall be 15 min to 30 min. For muscle samples that are less digestible, the digestion time should be increased but, unless otherwise validated for a particular sample matrix, the time shall not exceed 45 min.

4.5 Filtration of the digest fluid

Following digestion, the digest fluid shall be filtered through a sieve with specific mesh (6.9). Retained larvae shall be rinsed with tap water.

4.6 Verification of findings

If positive or doubtful findings occur, confirmation and identification at the species level should be performed by a qualified reference laboratory, by means of morphological and/or molecular methods.

1) Smasher and Stomacher® are examples of suitable products available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of these products.

5 Reagents

5.1 **Tap water.**

5.2 **Hydrochloric acid, 25 %.**

5.3 **Pepsin**, powder or granular: 1:10 000 NF, 1:12 500 BP, 2 000 FIP; liquid: 660 U/ml.

Pepsin is commercially available in powder, granular and liquid forms. The activity of the pepsin shall be certified. The pepsin shall be stored according to the manufacturer's recommendation.

The use of a liquid pepsin formulation can be advantageous as it could reduce the risk of occupational hazard, such as allergic reaction in laboratory staff.

NOTE The activity of pepsin powder is expressed per gram, either in "NF" (US National Formulary), "BP" (British Pharmacopoeia) or "FIP" (Fédération Internationale de Pharmacie). The activity of liquid pepsin is expressed in European Pharmacopoeia units per millilitre with a minimum of 660 U/ml. Other pepsin activities can be used, provided the final activity in the digest fluid is equivalent to the activity of 10 g of 1:10 000 NF.

5.4 **90 % ethanol.**

5.5 **Acidic solution, 1 % acetic acid.**

6 Equipment and consumables

Usual microbiological laboratory equipment in accordance with ISO 7218 and, in particular, the following.

6.1 **Labelled collection trays or plastic bags**, for samples.

6.2 **Knives, scissors and forceps**, for cutting samples.

6.3 **Calibrated scale**, for weighing samples and/or pepsin, accuracy $\pm 0,1$ g.

6.4 **Magnetic stirrer**, with an adjustable heating plate or magnetic stirrer put in an incubator.

6.5 **Polytetrafluoroethylene (PTFE)-coated triangular stir bar**, 5 cm in length, minimum.

6.6 **Thermometer**, accurate to at least $\pm 0,5$ °C, minimum range of temperatures from 20 °C to 70 °C.

6.7 **Glass beakers**, appropriate capacity.

6.8 **Aluminium foil or lids**, to cover the top of the glass beaker.

6.9 **Sieve**, made of brass or stainless steel, mesh size around 180 microns to 500 microns, with diameter of approximately 10 cm or larger.

6.10 **Petri dishes**, approximately 90 mm in diameter, for larval counting.

6.11 **Small vials**, for the collection of recovered larvae.

6.12 **Stereo-microscope**, with a sub-stage transmitted adjustable light source.

7 Sampling

Sampling is not part of the method specified in this document, but it is indicated in [Annex A](#). If there are no specific International Standards dealing with sampling of fish, it is recommended that the interested parties come to an agreement on this subject.

8 Procedure

WARNING — Laboratory staff performing the procedure shall wear disposable gloves, a mask and a laboratory coat. When handling pepsin, contact with the powder and inhalation of dusts (mask) should be avoided. The use of a liquid pepsin formulation can be advantageous as it could reduce the risk of an allergic reaction in laboratory staff.

8.1 Preparation of the sample

Samples used for testing can be fresh, frozen or lightly processed fish, and internal organs (e.g. liver, ovary). Their condition shall be reported before testing.

Each sample shall be weighed and its weight shall be reported. The minimum individual sample size for testing by digestion shall be at least 25 g and no more than 200 g.

Whole fish shall be manually eviscerated, skinned and filleted, and the viscera and each flesh side/fillet can be put in separate transparent plastic bags to be analysed independently, depending on fillet size.

Manually ease apart the sample or, alternatively, use a Smasher/Stomacher²⁾.

If any larvae are found before digestion, they should be properly preserved for further observation, as described in [8.5](#).

For lightly processed fish containing, for example, oil and spices, the sample shall be rinsed with tap water before digestion, to remove such ingredients as much as possible.

8.2 Preparation of the digest fluid

The pepsin used for the preparation of digest fluid shall have the appropriate activity required for digestion ([5.3](#)). Digest fluid shall be made fresh for each analysis. The steps for preparing digest fluid are as follows:

- a) add 16 ml of hydrochloric acid (see [Clause 5](#)) to a glass beaker containing 2 l of tap water preheated to $35\text{ °C} \pm 2\text{ °C}$;
- b) place a stirring rod in the beaker, place the beaker on a preheated magnetic stirrer or on a magnetic stirrer in the incubator, and commence the stirring;
- c) add 10 g of powder or granular pepsin (1:10 000 NF) or 30 ml of liquid pepsin (660 U/ml).

NOTE 1 The most critical step is the obligatory sequence of mixing of the digest fluid: a) water, b) hydrochloric acid, and c) pepsin. This will prevent degradation of pepsin that could result from direct exposure to concentrated hydrochloric acid.

NOTE 2 Stock solutions of hydrochloric acid are available in formulations other than 25 % and are adjusted accordingly. For example, if a 37 % stock solution of hydrochloric acid (molar concentration: 12,1) is used, 11 ml is added to 2 l of preheated tap water.

NOTE 3 Other activities of pepsin can be used, provided the final activity in the digest fluid is equivalent to the activity of 10 g of 1:10 000 NF ([5.3](#)).

2) Smasher and Stomacher® are examples of suitable products available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of these products.

8.3 Digestion of the sample in the glass beaker

- a) After preparation, the sample is transferred to a 2 l glass beaker, which contains the digest fluid (see 8.2).
- b) Cover the glass beaker with aluminium foil to keep a constant temperature of $37\text{ °C} \pm 2\text{ °C}$, and reduce evaporation.
- c) Regularly monitor the temperature of the digest fluid using a thermometer.
- d) Ensure that, during stirring, the digest fluid rotates at a sufficiently high speed to create a deep vortex without splashing.
- e) Digest the sample for 15 min to 30 min. If necessary, the digestion time can be increased but it shall not exceed 45 min in total.

NOTE The optimum temperature range of pepsin is 37 °C to 40 °C . However, under such conditions, nematodes belonging to the genus *Hysterothylacium* (considered to be non-zoonotic) will be killed and partially digested, which could negatively affect proper morphological or molecular identification.

8.4 Filtration of the digest fluid

Switch off the stirrer and immediately pour carefully (to avoid overflow) the digestion solution through the sieve into a beaker. Anisakidae larvae can be detected on the sieve.

The digestion process is considered satisfactory if residual debris remaining on the sieve consists primarily of indigestible tissue of no greater than 5 % of the original sample mass. If undigested tissue in excess remains on the sieve, the digestion procedure shall be repeated. In the case of excessive undigested tissue, a new sample shall be collected and the entire method repeated (i.e. start procedure again from 8.3).

NOTE 1 For frozen samples, which have been stored for several months at -20 °C or at lower temperatures, the larvae can appear damaged and/or fragmented and molecular identification can be difficult or impossible.

NOTE 2 In the case of fragments, a microscopic distinction can be made between front and rear ends. If fragments occur, it is documented whether they are front or rear ends. From this information, it is possible to estimate the minimum and maximum number of larvae.

8.5 Microscopic examination

Anisakidae larvae shall be collected from the sieve^[6] and examined under the stereomicroscope with transmitted light at $10\times$ to $20\times$ magnification for their morphological identification or processed for further analyses.

The larvae can be transferred in a vial filled with 90 % ethanol (see 5.4) and stored at a temperature range between -20 °C and 10 °C up to five years.

NOTE Viable larvae show active movements. If larval motility is not observed, larvae can be kept for 24 h in acidic solution ($\text{pH} < 4$) (see 5.5) at room temperature to assess motility.

9 Expression of the results

Results shall be expressed as “present” or “absent” and the number of Anisakidae larvae in “x” grams of sample, per fillet section.

NOTE If requested or appropriate, the localization of larvae detected, as well as their viability, can be reported.

If doubtful findings occur, confirmation and identification at the species level by molecular methods should be performed by a qualified reference laboratory.

10 Performance characteristics of the method

The values of these characteristics have been determined by a ring trial on the method organized within the framework of a European project^[3].

The accuracy, defined as the closeness of a measurement to the true value, is 98 % (95 % CI: 82,8 % to 99,1 %).

The sensitivity, defined as the percentage of samples correctly identified as positives, is 96 % (95 % CI: 79,6 % to 99,9 %).

The specificity, the percentage of samples correctly identified as negatives, is 100 % (95 % CI: 47,8 % to 100 %).

The limit of detection of the method is 1 larva in 100 g sample.

11 Test report

A laboratory worksheet (see [Annex B](#)) should be used by analysts to record data for test reports and is, therefore, a critical document for quality audits and trace-back investigations. Key components of the laboratory worksheet include sample tracking information, documentation that the method has been performed correctly by qualified personnel, equipment used (magnetic stirrer or incubator, thermometer), reagents used (batch number, date of preparation), documentation of problems and irregularities, and a written record of results. Laboratory worksheets should be stored according to the requirements of the competent authority.

12 Quality assurance

Each laboratory shall have a system of adequate documentation that demonstrates that anisakid testing was correctly performed according to appropriate quality assurance standards. There are no internal quality controls that can be used while performing the method.