
Oil of black pepper (*Piper nigrum* L.)

Huile essentielle de poivre noir (*Piper nigrum* L.)

STANDARDSISO.COM : Click to view the full PDF of ISO 3061:2008



PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

STANDARDSISO.COM : Click to view the full PDF of ISO 3061:2008



COPYRIGHT PROTECTED DOCUMENT

© ISO 2008

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 3061 was prepared by Technical Committee ISO/TC 54, *Essential oils*.

This second edition cancels and replaces the first edition (ISO 3061:1979), which has been technically revised.

STANDARDSISO.COM : Click to view the full PDF of ISO 3061:2008

[STANDARDSISO.COM](https://standardsiso.com) : Click to view the full PDF of ISO 3061:2008

Oil of black pepper (*Piper nigrum* L.)

1 Scope

This International Standard specifies certain characteristics of oil of black pepper (*Piper nigrum* L.), with a view to facilitating the assessment of its quality.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO/TR 210, *Essential oils — General rules for packaging, conditioning and storage*

ISO/TR 211, *Essential oils — General rules for labelling and marking of containers*

ISO 212, *Essential oils — Sampling*

ISO 279, *Essential oils — Determination of relative density at 20 °C — Reference method*

ISO 280, *Essential oils — Determination of refractive index*

ISO 592, *Essential oils — Determination of optical rotation*

ISO 875, *Essential oils — Evaluation of miscibility in ethanol*

ISO 11024-1, *Essential oils — General guidance on chromatographic profiles — Part 1: Preparation of chromatographic profiles for presentation in standards*

ISO 11024-2, *Essential oils — General guidance on chromatographic profiles — Part 2: Utilization of chromatographic profiles of samples of essential oils*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

essential oil of black pepper

essential oil obtained by steam distillation of the whole or broken unripe fruits of *Piper nigrum* L. of the Piperaceae family.

NOTE For information on the CAS number, see ISO/TR 21092^[2].

4 Requirements

4.1 Appearance

Clear mobile liquid.

4.2 Colour

Colourless or light coloured (yellow, green, blue).

4.3 Odour

Spicy, woody.

4.4 Relative density at 20 °C, d_{20}^{20}

	India	Sri Lanka	Indonesia	Madagascar
Minimum	0,864	0,861	0,861	0,864
Maximum	0,880	0,876	0,885	0,884

4.5 Refractive index at 20 °C

	India	Sri Lanka	Indonesia	Madagascar
Minimum	1,478	1,475	1,480	1,475
Maximum	1,487	1,490	1,493	1,490

4.6 Optical rotation at 20 °C

	India	Sri Lanka	Indonesia	Madagascar
Minimum	-18°	-17°	-17°	-18°
Maximum	-7°	-8°	-6°	+20°

4.7 Miscibility with 95 % (volume fraction) ethanol at 20 °C

It shall not be necessary to use more than 3 volumes of 95 % (volume fraction) ethanol to obtain a clear solution with 1 volume of essential oil.

4.8 Chromatographic profile

Analysis of the essential oil shall be carried out by gas chromatography. In the chromatogram obtained, the representative and characteristics components shown in Table 1 shall be identified. The percentages of these components, indicated by the integrator, shall be as shown in Table 1. This constitutes the chromatographic profile of the essential oil.

Table 1 — Chromatographic profile

Components	India		Sri Lanka		Indonesia		Madagascar	
	Min.	Max.	Min.	Max.	Min.	Max.	Min.	Max.
α -Pinene	3	12	10	16	2,5	10	8	26
β -Pinene	5	12	9	12	nd ^a	3	7	15
Sabinene	6	15	10	17	4	10	nd ^a	4
3-Carene	3	15	5	11	9	20	7	16
Limonene	10	17	13	16	7	12	10	25
δ -Elemene	0,5	3,5	0,5	3	0,5	3	0,5	4,5
α -Copaene	0,5	4,5	1,5	4	0,2	1	nd ^a	0,5
β -Caryophyllene	12	29	12	21	25	40	10	25
Germacrene D	nd ^a	2	nd ^a	1	nd ^a	1	1	6,5
β -Selinene	0,5	3,5	nd ^a	2	3	6	1	4,5
α -Selinene	nd ^a	3	0,5	2	2,5	5	0,5	3
Caryophyllene oxide	nd ^a	1						

^a non-detectable.

NOTE The chromatographic profile is normative, contrary to typical chromatograms given for information in Annex A.

4.9 Flashpoint

Information on the flashpoint is given in Annex B.

5 Sampling

See ISO 212.

Minimum volume of the final sample: 25 ml.

NOTE This volume is sufficient for each of the tests specified in this International Standard to be carried out at least once.

6 Test methods

6.1 Relative density at 20 °C, d_{20}^{20}

See ISO 279.

6.2 Refractive index at 20 °C

See ISO 280.

6.3 Optical rotation at 20 °C

See ISO 592.

6.4 Miscibility with 95 % (volume fraction) ethanol at 20 °C

See ISO 875.

6.5 Chromatographic profile

See ISO 11024-1 and ISO 11024-2.


7 Packing, labelling, marking and storage

See ISO/TR 210 and ISO/TR 211.

STANDARDSISO.COM : Click to view the full PDF of ISO 3061:2008

Annex A
(informative)

Typical chromatogram of the analysis by gas chromatography of the essential oil of black pepper (*Piper nigrum* L.)



Peak identification


- 1 α -Thujene
- 2 α -Pinene
- 3 Sabinene
- 4 β -Pinene
- 5 Myrcene
- 6 α -Phellandrene
- 7 3-Carene
- 8 Limonene + 1,8-Cineole
- 9 δ -Elemene
- 10 α -Copaene
- 11 β -Caryophyllene
- 12 α -Humulene
- 13 Germacrene D
- 14 β -Selinene
- 15 α -Selinene
- 16 Caryophyllene oxide

Operating conditions

Column: silica capillary; length 50 m; internal diameter 0,2 mm
 Stationary phase: polydimethylsiloxane [HP-1¹]
 Film thickness: 0,25 μ m
 Oven temperature: 65 °C to 230 °C, at a rate of 2 °C/min
 Injector temperature: 230 °C
 Detector temperature: 250 °C
 Detector: flame ionization type
 Carrier gas: hydrogen
 Volume injected: 0,2 μ l
 Carrier gas flow rate: 1 ml/min
 Split ratio: 1/100

Figure A.1 — Typical chromatogram taken on an apolar column, Madagascar type

1) Example of a suitable product available commercially. This information is given for the convenience of users of this International Standard, and does not constitute an endorsement of this product by ISO. Equivalent products may be used if they can be shown to lead to the same results.

**Peak identification**


- 1 α -Pinene + α -Thujene
- 2 β -Pinene
- 3 Sabinene
- 4 3-Carene
- 5 α -Phellandrene + Myrcene
- 6 Limonene
- 7 *p*-Cymene
- 8 Terpinolene
- 9 δ -Elemene
- 10 α -Copaene
- 11 β -Caryophyllene + Terpinen-4-ol
- 12 α -Humulene
- 13 Germacrene D
- 14 β -Selinene
- 15 α -Selinene
- 16 Caryophyllene oxide

Operating conditions

Column: silica capillary; length 50 m; internal diameter 0,2 mm
 Stationary phase: poly(ethylene glycol) [Carbowax²]
 Film thickness: 0,25 μ m
 Oven temperature: 65 °C to 230 °C, at a rate of 10 °C/min
 Injector temperature: 230 °C
 Detector temperature: 250 °C
 Detector: flame ionization type
 Carrier gas: hydrogen
 Volume injected: 0,2 μ l
 Carrier gas flow rate: 1,1 ml/min
 Split ratio: 1/100

Figure A.2 — Typical chromatogram taken on a polar column, Madagascar type

2) Example of a suitable product available commercially. This information is given for the convenience of users of this International Standard, and does not constitute an endorsement of this product by ISO. Equivalent products may be used if they can be shown to lead to the same results.

**Peak identification**


- 1 α -Thujene
- 2 α -Pinene
- 3 Sabinene
- 4 β -Pinene
- 5 Myrcene
- 6 α -Phellandrene
- 7 3-Carene
- 8 Limonene + 1,8-Cineole
- 9 δ -Elemene
- 10 α -Copaene
- 11 β -Caryophyllene
- 12 α -Humulene
- 13 Germacrene D
- 14 β -Selinene
- 15 α -Selinene
- 16 Caryophyllene oxide

Operating conditions

Column: silica capillary; length 30 m; internal diameter 0,25 mm
 Stationary phase: polydimethylsiloxane [HP-1³]
 Film thickness: 0,25 μ m
 Oven temperature: isothermal at 60 °C for 1 min, then programming temperature from 60 °C to 240 °C at a rate of 2 °C/min, then isothermal at 240 °C for 10 min
 Injector temperature: 250 °C
 Detector temperature: 300 °C
 Detector: flame ionization type
 Carrier gas: hydrogen
 Volume injected: 0,5 μ l
 Carrier gas flow rate: 100 ml/min
 Split ratio: 1/100

Figure A.3 — Typical chromatogram taken on an apolar column, Indian type

3) Example of a suitable product available commercially. This information is given for the convenience of users of this International Standard, and does not constitute an endorsement of this product by ISO. Equivalent products may be used if they can be shown to lead to the same results.

**Peak identification**

- 1 α -Pinene + α -Thujene
- 2 β -Pinene
- 3 Sabinene
- 4 3-Carene
- 5 α -Phellandrene + Myrcene
- 6 Limonene
- 7 *p*-Cymene
- 8 Terpinolene
- 9 δ -Elemene
- 10 α -Copaene
- 11 β -Caryophyllene + Terpinen-4-ol
- 12 α -Humulene
- 13 Germacrene D
- 14 β -Selinene
- 15 α -Selinene
- 16 Caryophyllene oxide

Operating conditions

Column: silica capillary; length 30 m; internal diameter 0,25 mm
 Stationary phase: poly(ethylene glycol) [Innowax⁴]
 Film thickness: 0,25 μ m
 Oven temperature: isothermal at 60 °C for 1 min, then programming temperature from 60 °C to 240 °C at a rate of 2 °C/min, then isothermal at 240 °C for 10 min
 Injector temperature: 250 °C
 Detector temperature: 300 °C
 Detector: flame ionization type
 Carrier gas: hydrogen
 Volume injected: 0,5 μ l
 Carrier gas flow rate: 100 ml/min
 Split ratio: 1/100

Figure A.4 — Typical chromatogram taken on a polar column, Indian type

4) Example of a suitable product available commercially. This information is given for the convenience of users of this International Standard, and does not constitute an endorsement of this product by ISO. Equivalent products may be used if they can be shown to lead to the same results.