

INTERNATIONAL STANDARD

ISO 3809

Third edition
2004-05-15

Oil of lime (cold pressed), Mexican type [*Citrus aurantifolia* (Christm.) Swingle], obtained by mechanical means

Huile essentielle de limette (exprimée à froid), type Mexique [Citrus aurantifolia (Christm.) Swingle], obtenue par procédés mécaniques

STANDARDSISO.COM : Click to view the full PDF of ISO 3809:2004



Reference number
ISO 3809:2004(E)

© ISO 2004

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

STANDARDSISO.COM : Click to view the full PDF of ISO 3809:2004

© ISO 2004

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 3809 was prepared by Technical Committee ISO/TC 54, *Essential oils*.

This third edition cancels and replaces the second edition (ISO 3809:1987), which has been technically revised.

STANDARDSISO.COM : Click to view the full PDF of ISO 3809:2004

Oil of lime (cold pressed), Mexican type [*Citrus aurantifolia* (Christm.) Swingle], obtained by mechanical means

1 Scope

This International Standard specifies certain characteristics of the oil of lime (cold pressed), Mexican type [*Citrus aurantifolia* (Christm.) Swingle], in order to facilitate assessment of its quality.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO/TR 210, *Essential oils — General rules for packaging, conditioning and storage*

ISO/TR 211, *Essential oils — General rules for labelling and marking of containers*

ISO 212, *Essential oils — Sampling*

ISO 279, *Essential oils — Determination of relative density at 20 °C — Reference method*

ISO 280, *Essential oils — Determination of refractive index*

ISO 592, *Essential oils — Determination of optical rotation*

ISO 1271, *Essential oils — Determination of carbonyl value — Free hydroxylamine method*

ISO 4715, *Essential oils — Quantitative evaluation of residue on evaporation*

ISO 4735, *Oils of citrus — Determination of CD value by ultraviolet spectrophotometric analysis*

ISO 11024-1, *Essential oils — General guidance on chromatographic profiles — Part 1: Preparation of chromatographic profiles for presentation in standards*

ISO 11024-2, *Essential oils — General guidance on chromatographic profiles — Part 2: Utilization of chromatographic profiles of samples of essential oils*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

oil of lime (cold pressed), Mexican type (type A)
essential oil obtained by centrifuging the emulsion of water, juice and oil obtained by crushing the whole fruits of *Citrus aurantifolia* (Christm.) Swingle, of the Rutaceae family

3.2

oil of lime (cold pressed), Mexican type (type B)
essential oil obtained by grating and/or squeezing the peel of fruits of *Citrus aurantifolia* (Christm.) Swingle, of the Rutaceae family, in the presence of water, followed by centrifuging the resulting emulsion of water and oil

NOTE 1 The principal areas of production are Mexico, the countries of Central America and the islands of the Caribbean.

NOTE 2 For information on the CAS number, see ISO/TR 21092.

4 Requirements

4.1 Appearance

Type A	Type B
Clear liquid, in which a waxy precipitate is usually present	

4.2 Colour

Type A	Type B
From yellowish green to green	Green to dark green

4.3 Odour

Type A	Type B
Characteristic of fresh lime peel. Fresh citrus lime-like odour	Fresh, with a note reminiscent of the pericarp. Fresh citrus lime odour with a pronounced juicy characteristic

4.4 Relative density at 20 °C, d_{20}^{20}

Type A		Type B	
min.	max.	min.	max.
0,875 0	0,884 0	0,880 0	0,888 0

4.5 Refractive index at 20 °C

Type A		Type B	
min.	max.	min.	max.
1,482 0	1,486 0	1,484 0	1,488 0

4.6 Optical rotation at 20 °C

Type A	Type B
Between +35,0° and +41,0°	This determination is often not possible because oils of this type are intensely coloured

4.7 Carbonyl value

Type A		Type B	
min.	max.	min.	max.
16 (corresponding to 4,5 % of carbonyl compounds expressed as citral)	31 (corresponding to 8,5 % of carbonyl compounds expressed as citral)	18 (corresponding to 5 % of carbonyl compounds expressed as citral)	35 (corresponding to 9,5 % of carbonyl compounds expressed as citral)

4.8 Residue on evaporation

Type A		Type B	
min.	max.	min.	max.
10,0 %	14,5 %	13,0 %	19,0 %

4.9 CD value

Type A	Type B
min.	min.
18,2	23,6

4.10 Chromatographic profile

Analysis of the essential oil shall be carried out by gas chromatography. In the chromatogram obtained, the representative and characteristic components shown in Tables 1 and 2, for type A and type B respectively, shall be identified. The proportions of these components, indicated by the integrator, shall be as shown in Tables 1 and 2. This constitutes the chromatographic profile of the essential oil.

4.11 Flashpoint

Information on the flashpoint is given in Annex B.

5 Sampling

See ISO 212.

Minimum volume of test sample: 25 ml.

NOTE This volume allows each of the tests specified in this International Standard to be carried out at least once.

6 Test methods

6.1 Relative density at 20 °C, d_{20}^{20}

See ISO 279.

6.2 Refractive index at 20 °C

See ISO 280.

6.3 Optical rotation at 20 °C

See ISO 592.

6.4 Carbonyl value

See ISO 1271.

Test portion: 5 g.

Standing time: 15 min.

6.5 Residue on evaporation

See ISO 4715.

Test portion: 5 g.

Evaporation time: 6 h.

6.6 CD value

See ISO 4735.

Point B: 370 nm approximately.

Maximum value: 312 nm to 315 nm approximately.

Point A: 280 nm approximately.

Dilution of 0,025 g of oil in 100 ml of 90 % ethanol (volume fraction).

Table 1 — Chromatographic profile (type A)

Component	Minimum %	Maximum %
α-Pinene	2,0	3,0
Sabinene	1,8	4,0
β-Pinene	18,0	24,0
Myrcene	1,0	2,0
p-Cymene	—	0,5
Limonene	42,0	50,0
γ-Terpinene	8,0	11,0
Terpinen-4-ol	0,2	0,6
α-Terpineol	0,2	0,6
n-Decanal	0,05	0,3
Neral	1,2	2,0
Geranial	2,0	3,0
Neryl acetate	0,1	0,35
Geranyl acetate	0,2	0,4
β-Caryophyllene	0,5	1,5
α-Bergamotene	1,0	1,9
α-Farnesene	0,75	1,75
β-Bisabolene	1,0	1,5

NOTE 1 The chromatographic profile is normative, contrary to typical chromatograms given for information in Annex A.

NOTE 2 Expressed oils of lime may contain furo-coumarins. The bergapten content should be less than 2 000 mg/kg.

6.7 Chromatographic profile

See ISO 11024-1 and ISO 11024-2.

7 Packaging, labelling, marking and storage

See ISO/TR 210 and ISO/TR 211.

Table 2 — Chromatographic profile (type B)


Component	Minimum %	Maximum %
α-Pinene	1,7	2,0
Sabinene	2,0	3,0
β-Pinene	17,0	19,0
Myrcene	1,4	1,8
Neryl acetate	0,0	0,25
γ-Terpinene	9,0	9,7
Terpinen-4-ol	0,2	0,6
p-Cymene	—	0,5
Limonene	38,0	44,0
α-Terpineol	0,3	0,6
n-Decanal	0,15	0,35
Neral	2,0	2,5
Geranial	3,0	3,7
Geranyl acetate	0,3	0,6
β-Caryophyllene	1,5	1,9
α-Bergamotene	0,5	0,7
β-Bisabolene	4,0	4,5

NOTE 1 The chromatographic profile is normative, contrary to typical chromatograms given for information in Annex A.

NOTE 2 Expressed oils of lime may contain furo-coumarins. The bergapten content should be less than 2 000 mg/kg.

Annex A (informative)

Typical chromatograms of the analysis by gas chromatography of the essential oil of lime (cold pressed), Mexican type [*Citrus aurantifolia* (Christm.) Swingle], obtained by mechanical means




Peak identification

1	α -Pinene	10	<i>n</i> -Decanal
2	Sabinene	11	Neral
3	β -Pinene	12	Geranial
4	Myrcene	13	Neryl acetate
5	<i>p</i> -Cymene	14	Geranyl acetate
6	Limonene	15	β -Caryophyllene
7	γ -Terpinene	16	α -Bergamotene
8	Terpinen-4-ol	17	α -Farnesene
9	α -Terpineol	18	β -Bisabolene

Operating conditions

Column: capillary; length 30 m; internal diameter 0,20 mm
 Stationary phase: poly(5 % diphenyl/95 % dimethyl siloxane) (SP-5[®])
 Film thickness: 20 μ m
 Oven temperature: isothermal at 75 °C for 5 min, then temperature programming from 75 °C to 100 °C at a rate of 5 °C/min, then from 100 °C to 220 °C at a rate of 6 °C/min, and isothermal at 220 °C for 8,5 min
 Injector temperature: 230 °C
 Detector temperature: 260 °C
 Detector: flame ionization type
 Carrier gas: helium
 Volume injected: 1 μ l
 Carrier gas flow rate: 206,84 kPa
 Split ratio: 1/100

Figure A.1 — Typical chromatogram taken on an apolar column (type A)


**Peak identification**

1	α -Pinene	8	α -Bergamotene
2	β -Pinene	9	β -Caryophyllene
3	Sabinene	10	Neral
4	Myrcene	11	α -Terpineol
5	Limonene	12	β -Bisabolene
6	γ -Terpinene	13	Neryl acetate
7	<i>p</i> -Cymene		

Operating conditions

Column: capillary; length 30 m; internal diameter 0,20 mm
 Stationary phase: poly(ethylene glycol) (Carbowax®)
 Film thickness: 20 μ m
 Oven temperature: isothermal at 75 °C for 5 min, then temperature programming from 75 °C to 100 °C at a rate of 5 °C/min, then from 100 °C to 220 °C at a rate of 6 °C/min and isothermal at 220 °C for 8,5 min
 Injector temperature: 230 °C
 Detector temperature: 260 °C
 Detector: flame ionization type
 Carrier gas: helium
 Volume injected: 1 μ l
 Carrier gas flow rate: 206,84 kPa
 Split ratio: 1/100

Figure A.2 — Typical chromatogram taken on a polar column (type A)


**Peak identification**

1	α -Pinene	10	<i>n</i> -Decanal
2	Sabinene	11	Neral
3	β -Pinene	12	Geranial
4	Myrcene	13	Neryl acetate
5	<i>p</i> -Cymene	14	Geranyl acetate
6	Limonene	15	β -Caryophyllene
7	γ -Terpinene	16	α -Bergamotene
8	Terpinen-4-ol	17	α -Farnesene
9	α -Terpineol	18	β -Bisabolene

Operating conditions

Column: capillary; length 30 m; internal diameter 0,20 mm
 Stationary phase: poly(5 % diphenyl/95 % dimethyl siloxane) (SP-5[®])
 Film thickness: 20 μ m
 Oven temperature: isothermal at 75 °C for 5 min, then temperature programming from 75 °C to 100 °C at a rate of 5 °C/min, then from 100 °C to 220 °C at a rate of 6 °C/min, and isothermal at 220 °C for 8,5 min
 Injector temperature: 230 °C
 Detector temperature: 260 °C
 Detector: flame ionization type
 Carrier gas: helium
 Volume injected: 1 μ l
 Carrier gas flow rate: 206,84 kPa
 Split ratio: 1/100

Figure A.3 — Typical chromatogram taken on an apolar column (type B)



Peak identification

- | | |
|-----------------------|--------------------------|
| 1 α -Pinene | 8 α -Bergamotene |
| 2 β -Pinene | 9 β -Caryophyllene |
| 3 Sabinene | 10 Neral |
| 4 Myrcene | 11 α -Terpineol |
| 5 Limonene | 12 β -Bisabolene |
| 6 γ -Terpinene | 13 Neryl acetate |
| 7 <i>p</i> -Cymene | |

Operating conditions

Column: capillary; length 30 m; internal diameter 0,20 mm
 Stationary phase: poly(ethylene glycol) (Carbowax®)
 Film thickness: 20 μ m
 Oven temperature: isothermal at 75 °C for 5 min, then temperature programming from 75 °C to 100 °C at a rate of 5 °C/min, then from 100 °C to 220 °C at a rate of 6 °C/min and isothermal at 220 °C for 8,5 min
 Injector temperature: 230 °C
 Detector temperature: 260 °C
 Detector: flame ionization type
 Carrier gas: helium
 Volume injected: 1 μ l
 Carrier gas flow rate: 206,84 kPa
 Split ratio: 1/100

Figure A.4 — Typical chromatogram taken on a polar column (type B)