
International Standard



5521

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION • МЕЖДУНАРОДНАЯ ОРГАНИЗАЦИЯ ПО СТАНДАРТИЗАЦИИ • ORGANISATION INTERNATIONALE DE NORMALISATION

Fruits, vegetables and derived products — Qualitative method for the detection of sulphur dioxide

Fruits, légumes et produits dérivés — Méthode qualitative de recherche du dioxyde de soufre

First edition — 1981-08-01

STANDARDSISO.COM : Click to view the full PDF of ISO 5521:1981

UDC 634/635 : 543.061 : 546.224-31

Ref. No. ISO 5521-1981 (E)

Descriptors : fruit and vegetable products, agricultural products, fruits, vegetables, qualitative analysis, detection, sulphur dioxide.

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards institutes (ISO member bodies). The work of developing International Standards is carried out through ISO technical committees. Every member body interested in a subject for which a technical committee has been set up has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work.

Draft International Standards adopted by the technical committees are circulated to the member bodies for approval before their acceptance as International Standards by the ISO Council.

International Standard ISO 5521 was developed by Technical Committee ISO/TC 34, *Agricultural food products*, and was circulated to the member bodies in June 1979.

It has been approved by the member bodies of the following countries :

Australia	Germany, F. R.	New Zealand
Austria	Hungary	Philippines
Brazil	India	Poland
Bulgaria	Israel	Portugal
Canada	Kenya	Romania
Cyprus	Libyan Arab Jamahiriya	South Africa, Rep. of
Czechoslovakia	Malaysia	Spain
Egypt, Arab Rep. of	Mexico	Thailand
France	Netherlands	Yugoslavia

No member body expressed disapproval of the document.

Fruits, vegetables and derived products — Qualitative method for the detection of sulphur dioxide

1 Scope and field of application

This International Standard specifies a qualitative method for the detection of free or bound sulphur dioxide in fruits, vegetables and derived products.

The method is applicable to products having sulphur dioxide contents of 2 mg or more per kilogram (or per litre).

For products containing garlic or onion, see clause 6.

2 Principle

2.1 Free sulphur dioxide

After acidification of the test portion, decolouration by the liberated gaseous sulphur dioxide of a starch indicator paper, previously coloured blue by a trace of iodine vapour.

2.2 Bound sulphur dioxide

Addition of alkali to the test portion, followed by the procedure described in 2.1.

3 Reagents

All reagents shall be of recognized analytical quality. The water used shall be distilled water or water of at least equivalent purity, recently boiled.

3.1 Starch, approximately 5 g/l solution.

During preparation, boil the solution for 10 min.

The solution will keep only for a few days.

3.2 Starch paper.

Dip a white filter paper, of close texture, into the starch solution (3.1) three times and dry in an oven at 30 °C after each impregnation. Cut into strips of 2 cm × 5 cm.

3.3 Potassium iodide, approximately 1 g/l solution (free from iodate).

3.4 Iodine, approximately 12 g/l solution.

3.5 Phosphoric acid, approximately 50 % (m/m) solution, $\rho_{20} = 1,34$ g/ml.

3.6 Sodium hydroxide, approximately 40 g/l solution.

4 Apparatus

Usual laboratory apparatus, not otherwise specified, and in particular :

4.1 Flasks, of clear glass, of capacity 150 ml, and having wide mouths and ground necks.

4.2 Ground glass stoppers, to fit the flasks (4.1).

5 Procedure

5.1 Preparation of starch indicator paper

Immediately before use, wet a strip of starch paper (3.2) with two drops of the potassium iodide solution (3.3) and attach it to the bottom of a glass stopper (4.2), for example by means of adhesive tape. Place a few millilitres of the iodine solution (3.4) in one of the flasks (4.1). Insert the stopper (the starch paper shall be suspended in the air space above the iodine). Leave for 5 to 10 s. A clearly visible pale blue colour will develop on the starch indicator paper which has to be used immediately.

5.2 Test portion

Place in another flask (4.1) about 20 ml of the product to be analysed (in the case of liquid products) or about 20 g of finely comminuted product (in the case of solid products) suspended in 20 ml of water.

5.3 Detection of free sulphur dioxide

5.3.1 Acidify the test portion with a few drops of the phosphoric acid solution (3.5).

5.3.2 Immediately close the flask by means of the stopper with the starch indicator paper, prepared as described in 5.1, attached. In the presence of traces of sulphur dioxide [2 mg per kilogram (or per litre)] the indicator paper will be decolorized in less than 5 min.