
**Rennet caseins and caseinates —
Determination of ash (Reference method)**

*Caséines présure et caséinates — Détermination des cendres
(Méthode de référence)*

STANDARDSISO.COM : Click to view the full PDF of ISO 5545:2008



Reference numbers
ISO 5545:2008(E)
IDF 90:2008(E)

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. Neither the ISO Central Secretariat nor the IDF accepts any liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies and IDF national committees. In the unlikely event that a problem relating to it is found, please inform the ISO Central Secretariat at the address given below.

STANDARDSISO.COM : Click to view the full PDF of ISO 5545:2008



COPYRIGHT PROTECTED DOCUMENT

© ISO and IDF 2008

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO or IDF at the respective address below.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

International Dairy Federation
Diamant Building • Boulevard Auguste Reyers 80 • B-1030 Brussels
Tel. + 32 2 733 98 88
Fax + 32 2 733 04 13
E-mail info@fil-idf.org
Web www.fil-idf.org

Published in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 5545|IDF 90 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*, and the International Dairy Federation (IDF). It is being published jointly by ISO and IDF.

This second edition of ISO 5545|IDF 90 cancels and replaces the first edition (ISO 5545:1978), of which it constitutes a minor revision.

STANDARDSISO.COM : Click to view the full PDF of ISO 5545:2008

Foreword

IDF (the International Dairy Federation) is a non-profit organization representing the dairy sector worldwide. IDF membership comprises National Committees in every member country as well as regional dairy associations having signed a formal agreement on cooperation with IDF. All members of IDF have the right to be represented at the IDF Standing Committees carrying out the technical work. IDF collaborates with ISO in the development of standard methods of analysis and sampling for milk and milk products.

Draft International Standards adopted by the Action Teams and Standing Committees are circulated to the National Committees for voting. Publication as an International Standard requires approval by at least 50 % of the IDF National Committees casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. IDF shall not be held responsible for identifying any or all such patent rights.

ISO 5545|IDF 90 was prepared by the International Dairy Federation (IDF) and Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*. It is being published jointly by IDF and ISO.

All work was carried out by the former Joint ISO-IDF-AOAC Group of Experts (E11-E701) which is now part of the Joint ISO-IDF Action Team on *Physical properties and rheological tests*, of the Standing Committee on *Minor components and characterization of physical properties*.

This edition of ISO 5545|IDF 90 cancels and replaces IDF 90:1979, of which it constitutes a minor revision.

STANDARDSISO.COM : Click to view the full PDF of ISO 5545:2008

Rennet caseins and caseinates — Determination of ash (Reference method)

1 Scope

This International Standard specifies a reference method for the determination of the ash of caseins obtained by rennet precipitation and of caseinates, with the exception of ammonium caseinate.

NOTE For the determination of ash ("fixed ash") of acid caseins, of ammonium caseinates, of their mixtures with rennet casein and with caseinates, and of caseins of unknown type, see ISO 5544 | IDF 89 [2].

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 3310-1, *Test sieves — Technical requirements and testing — Part 1: Test sieves of metal wire cloth*

ISO 5550 | IDF 78, *Caseins and caseinates — Determination of moisture content (Reference method)*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

ash of rennet caseins or of caseinates

substances determined by the procedure described in this International Standard

NOTE Ash of rennet caseins or of caseinates is conventionally expressed as a percentage by mass, numerically equivalent to grams per 100 g of product.

4 Principle

A test portion is incinerated at (825 ± 25) °C. The residue is then weighed.

5 Apparatus

Usual laboratory apparatus, and in particular the following.

5.1 Analytical balance, capable of weighing to the nearest 0,000 1 g.

5.2 Silica or platinum dishes, of diameter about 70 mm and of depth 25 mm to 50 mm.

5.3 Electrical furnace, with air circulation, capable of being maintained at $(825 \pm 25) ^\circ\text{C}$.

5.4 Desiccator, containing an effective desiccant.

5.5 Grinding device, for grinding the laboratory sample, if necessary (see 7.1.4), without development of undue heat and without loss or absorption of moisture. A hammer-mill shall not be used.

5.6 Test sieve, wire cloth, diameter 200 mm, nominal size of aperture 500 μm , with receiver, complying with ISO 3310-1.

6 Sampling

A representative sample should have been sent to the laboratory. It should not have been damaged or changed during transport or storage.

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 707|IDF 50 [1].

7 Procedure

7.1 Preparation of the test sample

7.1.1 Thoroughly mix the laboratory sample by repeatedly shaking and inverting the container (if necessary, after having transferred all of the laboratory sample to an airtight container of sufficient capacity to allow this operation to be carried out).

7.1.2 Transfer about 50 g of the thoroughly mixed laboratory sample to the test sieve (5.6).

7.1.3 If the 50 g portion directly passes or almost completely passes the sieve, use for the determination the sample as prepared in 7.1.1.

7.1.4 Otherwise, grind the 50 g portion, using the grinding device (5.5), until it passes the sieve. Immediately transfer all of the sieved sample to an airtight container of sufficient capacity and mix thoroughly by repeatedly shaking and inverting. During these operations, take precautions to avoid any change in the water content of the product.

7.1.5 After the test sample has been prepared, the determination (7.4) should proceed as soon as possible.

Clean the device after grinding each sample.

7.2 Preparation of the dish

Heat the dish (5.2) in the electrical furnace (5.3), maintained at $825 ^\circ\text{C}$, for 30 min. Allow the dish to cool in the desiccator (5.4) to the temperature of the balance room and weigh to the nearest 0,1 mg.

7.3 Test portion

Weigh, to the nearest 0,1 mg, directly in or by difference approximately 3,000 0 g of the test sample (7.1) into the prepared dishes.

7.4 Determination

Heat the dish containing the test portion on a low flame until the test portion is completely charred, taking care that it does not burst into flame.

Transfer the dish to the electrical furnace (5.3), maintained at $(825 \pm 25) ^\circ\text{C}$, and heat for at least 1 h until all carbon has disappeared from the dish. Allow the dish to cool in the desiccator (5.4) to the temperature of the balance room and weigh to the nearest 0,1 mg.

Repeat the operations of heating in the electrical furnace (5.3), cooling and weighing, until the mass remains constant to within 1 mg or begins to increase. Record the minimum mass.

8 Expression of results

8.1 Calculation

8.1.1 The ash of the sample, w_a , as a percentage by mass, is given by Equation (1):

$$w_a = \frac{m_1 - m_2}{m_0} \times 100 \quad (1)$$

where

m_0 is the mass, in grams, of the test portion (7.3);

m_1 is the mass, in grams, of dish and residue (7.4);

m_2 is the mass, in grams, of the prepared dish (7.2).

Calculate the ash to the nearest 0,01 % by mass and report the final result to the nearest 0,1 % by mass.

8.1.2 To calculate the ash of the sample on the dry basis, as a percentage by mass, multiply the result obtained from Equation (1) by Factor (2):

$$\frac{100}{100 - w_w} \quad (2)$$

where w_w is the water content, as a percentage by mass, of the sample determined according to ISO 5550 | IDF 78.

8.2 Precision

8.2.1 Repeatability

The absolute difference between two independent single test results, obtained using the same method on identical test material in the same laboratory by the same operator using the same equipment within a short interval of time will in not more than 5 % of cases be greater than 0,15 g of ash per 100 g of product (numerically equivalent to 0,15 % by mass).

8.2.2 Reproducibility

The absolute difference between two independent single test results, obtained using the same method on identical test material in different laboratories with different operators using different equipment, will in not more than 5 % of cases be greater than 0,25 g of ash per 100 g of product (numerically equivalent to 0,25 % by mass).

9 Test report

The test report shall specify:

- a) all the information required for the complete identification of the sample;
- b) the sampling method used, if known;
- c) the test method used, together with a reference to this International Standard;
- d) all operating details not specified in this International Standard, or regarded as optional, together with details of any incident that may have influenced the result(s);
- e) the test result(s) obtained, and if the repeatability has been checked, the final quoted results obtained.

STANDARDSISO.COM : Click to view the full PDF of ISO 5545:2008