

# International Standard



# 6651

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION • МЕЖДУНАРОДНАЯ ОРГАНИЗАЦИЯ ПО СТАНДАРТИЗАЦИИ • ORGANISATION INTERNATIONALE DE NORMALISATION

## Animal feeding stuffs — Determination of aflatoxin B<sub>1</sub> content

*Aliments des animaux — Dosage de l'aflatoxine B<sub>1</sub>*

First edition — 1983-04-01

STANDARDSISO.COM : Click to view the full PDF of ISO 6651:1983

UDC 636.085 : 543.544

Ref. No. ISO 6651-1983 (E)

Descriptors : animal feeding products, chemical analysis, determination of content, aflatoxins.

Price based on 8 pages

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of developing International Standards is carried out through ISO technical committees. Every member body interested in a subject for which a technical committee has been authorized has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work.

Draft International Standards adopted by the technical committees are circulated to the member bodies for approval before their acceptance as International Standards by the ISO Council.

International Standard ISO 6651 was developed by Technical Committee ISO/TC 34, *Agricultural food products*, and was circulated to the member bodies in March 1981.

It has been approved by the member bodies of the following countries:

Australia	Iran	Portugal
Austria	Iraq	Romania
Brazil	Ireland	South Africa, Rep. of
Canada	Israel	Spain
Chile	Italy	Sri Lanka
Denmark	Kenya	Tanzania
Dominican Republic	Korea, Rep. of	Thailand
Egypt, Arab Rep. of	Malaysia	Turkey
Ethiopia	Mexico	USSR
France	New Zealand	Yugoslavia
Hungary	Philippines	
India	Poland	

The member bodies of the following countries expressed disapproval of the document on technical grounds:

United Kingdom  
USA

This International Standard has also been approved by the International Union of Pure and Applied Chemistry (IUPAC).

# Animal feeding stuffs — Determination of aflatoxin B<sub>1</sub> content

## 1 Scope

This International Standard specifies two methods for the determination of the aflatoxin B<sub>1</sub> content of animal feeding stuffs.

## 2 Field of application

**2.1 Method A** is applicable to the following simple feeding stuffs :

- oilseeds and oilseed residues, and in particular groundnut, copra, linseed, soya, sesame, babassu palm;
- manioc meal;
- maize germ expeller;
- cereals and cereal products;
- pea meal;
- potato pulp and flour.

In the presence of substances interfering with the determination by method A, it is recommended that the determination be carried out in accordance with method B.

**2.2 Method B** is applicable to mixed feeding stuffs and to simple feeding stuffs not mentioned in 2.1.

This method is not applicable to feeding stuffs containing citrus pulp.

**2.3** The lower limit of detection of aflatoxin B<sub>1</sub> is 0,01 mg/kg.

## 3 Reference

ISO 6498, *Animal feeding stuffs — Preparation of test samples.*<sup>1)</sup>

## 4 Principle

Extraction of the test portion with chloroform, filtration, and purification of an aliquot portion on a silica gel column.

1) At present at the stage of draft.

2) The solvents should be used in covered tanks. When saturated tanks are specified, this is achieved by lining the tanks with absorbent paper and allowing the interiors to become saturated with solvent vapour.

Evaporation of the eluate and dissolution of the residue in a specified volume of chloroform or mixture of benzene and acetonitrile.

Thin-layer chromatography, one-dimensional for method A and two-dimensional for method B, of an aliquot portion of this solution.

Determination of the aflatoxin B<sub>1</sub> content, either visually or by fluorodensitometry, by examination of the chromatogram under UV light and comparison with known quantities of standard aflatoxin B<sub>1</sub> applied to the same plate as the test portion extract.

Confirmation of the identity of aflatoxin B<sub>1</sub> by formation of the hemiacetal derivative.

## 5 Reagents

All reagents shall be of recognized analytical quality. The water used shall be distilled water or water of at least equivalent purity.

**5.1 Chloroform**, stabilized with 0,5 to 1,0 % of 96 % (V/V) ethanol.

**5.2 n-hexane.**

**5.3 Diethyl ether**, anhydrous, free from peroxides.

**5.4 Benzene/acetonitrile (98 + 2) mixture.**

Mix 98 volumes of benzene with 2 volumes of acetonitrile.

**5.5 Chloroform/methanol (97 + 3) mixture.**

Mix 97 volumes of chloroform with 3 volumes of methanol.

**5.6 Developing solvents.**<sup>2)</sup>

**5.6.1 Chloroform/acetone (90 + 10) mixture.**

Mix 90 volumes of chloroform with 10 volumes of acetone, in an unsaturated tank.

**5.6.2 Diethyl ether/methanol/water (96 + 3 + 1) mixture.**

Mix 96 volumes of diethyl ether, 3 volumes of methanol and 1 volume of water, in an unsaturated tank.

**5.6.3 Diethyl ether/methanol/water (94 + 4,5 + 1,5) mixture.**

Mix 94 volumes of diethyl ether with 4,5 volumes of methanol and 1,5 volumes of water, in a saturated tank.

**5.6.4 Chloroform/methanol (94 + 6) mixture.**

Mix 94 volumes of chloroform with 6 volumes of methanol, in a saturated tank.

**5.6.5 Chloroform/methanol (97 + 3) mixture.**

Mix 97 volumes of chloroform with 3 volumes of methanol, in a saturated tank.

**5.7 Silica gel**, for column chromatography, of particle size 0,05 to 0,20 mm.

**5.8 Silica gel**, G-HR or equivalent, for thin layer chromatography.

**5.9 Diatomaceous earth** (Hyflosupercel), acid-washed.

**5.10 Sodium sulphate**, anhydrous granules.

**5.11 Trifluoroacetic acid.**

**5.12 Inert gas**, for example nitrogen.

**5.13 Sulphuric acid**, 50 % (V/V) solution.

**5.14 Aflatoxin B<sub>1</sub>**, standard solution containing about 0,1 µg of aflatoxin B<sub>1</sub> per millilitre, in the chloroform (5.1) or in the benzene/acetonitrile mixture (5.4).

**WARNING — Aflatoxins are highly carcinogenic and must be handled with great care.**

Prepare and check the solution as follows.

**5.14.1 Preparation of stock solution and determination of concentration**

Prepare a solution of aflatoxin B<sub>1</sub> in the chloroform (5.1) or the benzene/acetonitrile mixture (5.4) such that the concentration is between 8 and 10 µg/ml. Determine the absorption spectrum between 330 and 370 nm by means of the spectrophotometer (6.9).

Measure the absorbance (*A*) at 363 nm in the case of the chloroform solution, or at 348 nm in the case of the benzene/acetonitrile mixture solution.

Calculate the concentration of aflatoxin B<sub>1</sub>, in micrograms per millilitre of solution, from the formulae :

a) for the chloroform solution :

$$\frac{312 \times A \times 1\,000}{20\,600}$$

b) for the solution in the benzene/acetonitrile mixture :

$$\frac{312 \times A \times 1\,000}{19\,800}$$

**5.14.2 Dilution**

Dilute the stock solution (5.14.1), as appropriate, away from daylight, to obtain a standard solution with a concentration of aflatoxin B<sub>1</sub> of about 0,1 µg/ml.

If kept in a refrigerator at 4 °C, this solution is stable for 2 weeks.

**5.14.3 Testing of chromatographic purity of the solution**

Onto a plate (6.7), apply a spot of 5 µl of the standard aflatoxin B<sub>1</sub> solution of concentration 8 to 10 µg/ml (5.14.1). Develop the chromatogram as indicated in 8.5.1. Under UV light, the chromatogram shall show only one spot and no fluorescence shall be perceptible in the original deposition zone.

**5.15 Aflatoxin B<sub>1</sub> and B<sub>2</sub>** (see the warning in 5.14), solutions for qualitative testing, containing about 0,1 µg of aflatoxin B<sub>1</sub> and B<sub>2</sub> per millilitre, in the chloroform (5.1) or in the benzene/acetonitrile mixture (5.4).

These concentrations are given as a guide. They shall be adjusted so as to obtain the same intensity of fluorescence for both aflatoxins (see 8.5.1).

**6 Apparatus**

Usual laboratory equipment, and in particular :

**6.1 Grinder/mixer.**

**6.2 Sieve**, of aperture size 1,0 mm (see ISO 565).

**6.3 Shaking apparatus or magnetic stirrer.**

**6.4 Chromatographic tubes**, made of glass (internal diameter 22 mm, length 300 mm), with a PTFE tap and a 250 ml reservoir, plugged at the bottom end with cotton or glass wool.

**6.5 Rotary vacuum evaporator**, with a 500 ml round-bottomed flask.

**6.6 Apparatus for thin layer chromatography (TLC)**, i.e. that necessary for the preparation of the plates (6.7) and application of spots (capillary pipettes or microsyringes), a developing tank, and spraying apparatus for applying the sulphuric acid (5.13) to the plates.

**6.7 Glass TLC plates**, 200 mm × 200 mm, prepared as follows (the quantities indicated are sufficient to cover five plates).

Place 30 g of the silica gel (5.8) in a conical flask, add 60 ml of water, stopper and shake for 1 min. Spread the suspension on the plates so as to obtain a uniform layer 0,25 mm thick. Leave to dry in the air and then store in a desiccator containing silica gel. At the time of use, activate the plates by keeping them in an oven at 110 °C for 1 h.

Ready-to-use plates are suitable if they give results similar to those obtained with the plates prepared as indicated above.

**6.8 Long-wavelength (360 nm) UV lamp.**

The intensity of irradiation shall make it possible for a spot of 1,0 ng of aflatoxin B<sub>1</sub> to be clearly distinguished on a TLC plate at a distance of 10 cm from the lamp.

**WARNING** — UV light is dangerous to the eyes. Protective goggles shall be worn.

**6.9 Spectrophotometer**, suitable for making measurements in the UV region of the spectrum.

**6.10 Fluorodensitometer** (optional).

**6.11 Fluted filter paper.**

**6.12 Graduated tube**, of capacity 10,0 ml, with a polyethylene stopper.

**6.13 Conical flask**, of capacity 500 ml, with a ground glass stopper.

**6.14 Pipette**, of capacity 50 ml.

**6.15 Balance.**

## 7 Sampling

Take the laboratory sample from the material to be sampled in accordance with the International Standard for the material concerned unless sampling for the determination of aflatoxin is excluded from its field of application. If no appropriate International Standard exists, agreement should be reached between the parties concerned, taking into account the characteristics of the material being sampled.

## 8 Procedure

### 8.1 Preparation of the test sample

**8.1.1** If the sample contains more than 5 % of fat, it shall be defatted with light petroleum before grinding.

In such cases, the analytical results shall be expressed in terms of the mass of the non-defatted sample.

**8.1.2** Grind the laboratory sample so that it completely passes through the sieve (6.2). Mix thoroughly. (See ISO 6498.)

### 8.2 Test portion

Weigh 50 g of the prepared test sample into the conical flask (6.13).

### 8.3 Extraction

Add to the test portion (8.2) 25 g of the diatomaceous earth (5.9), 25 ml of water, and 250 ml of the chloroform (5.1) accurately measured from a measuring cylinder. Stopper the flask, and shake or stir for 30 min using the apparatus (6.3). Filter through the fluted filter paper (6.11), taking care to discard the first 10 ml of the filtrate, and subsequently collect at least 50 ml of the filtrate.

### 8.4 Column clean-up

#### 8.4.1 Preparation of the column

Fill two-thirds of the chromatographic tube (6.4) with the chloroform (5.1) and add 5 g of the sodium sulphate (5.10). Check that the upper surface of the sodium sulphate layer is flat, then add 10 g, in small portions, of the silica gel (5.7). Stir carefully after each addition to eliminate air bubbles. Leave to stand for 15 min and then carefully add 10 g of the sodium sulphate (5.10). Open the tap and allow the liquid to flow until it is just above the upper surface of the sodium sulphate layer. Close the tap.

#### 8.4.2 Purification

Transfer, by means of the pipette (6.14), 50 ml of the filtrate collected in 8.3 to a 250 ml conical flask, and add 100 ml of the *n*-hexane (5.2). Mix and quantitatively transfer the mixture to the column, rinsing the flask with the *n*-hexane. Open the tap and allow the liquid to flow at a rate of 8 to 12 ml/min until it is level with the upper surface of the sodium sulphate layer. Close the tap. Discard the liquid collected and pour 100 ml of the diethyl ether (5.3) into the column. Again open the tap and allow the liquid to flow until it is level with the upper surface of the sodium sulphate layer. During these operations, ensure that the column does not run dry.

Elute with 150 ml of the chloroform/methanol mixture (5.5) and collect the whole of the eluate in the 500 ml flask of the rotary evaporator (6.5). Evaporate to dryness on the rotary evaporator, preferably under a stream of inert gas (5.12), at a temperature not exceeding 50 °C, and under reduced pressure.

NOTE — If a rotary evaporator is not available, add a boiling aid and evaporate almost to dryness on a water bath.

Quantitatively transfer the residue, using the chloroform (5.1) or the benzene/acetonitrile mixture (5.4), to the 10 ml graduated tube (6.12). Again evaporate the solution, for example in a water bath, preferably under a stream of inert gas (5.12), and adjust the volume to 2,0 ml with the chloroform (5.1) or the benzene/acetonitrile mixture (5.4).

## 8.5 Thin-layer chromatography

### 8.5.1 Method A — One-dimensional thin-layer chromatography

#### 8.5.1.1 Choice of solvent

The choice of solvent (5.6.1, 5.6.2, 5.6.3, 5.6.4 or 5.6.5) shall be made beforehand to ensure that aflatoxins B<sub>1</sub> and B<sub>2</sub> are completely separated when the plate is developed, which depends on the batch of plates in use.

Place 25 µl of the qualitative solution (5.15) on the prepared plates (6.7) (one plate for each solvent to be checked). Follow the procedure in 8.5.1.2 for development, evaporation and irradiation. Two distinct spots are produced by a suitable solvent.

#### 8.5.1.2 Procedure

Onto a TLC plate (6.7), and using a capillary pipette or microsyringe, apply 20 mm from the lower edge, and at intervals of 20 mm, the volumes indicated below of the standard aflatoxin B<sub>1</sub> solution and the extract :

- 10, 15, 20, 30 and 40 µl of the standard aflatoxin B<sub>1</sub> solution (5.14);
- 10 µl of the extract obtained in 8.4.2 and, superimposed on the same point, 20 µl of the standard aflatoxin B<sub>1</sub> solution (5.14);
- 10 and 20 µl of the extract obtained in 8.4.2.

Develop the chromatogram in the dark using the developing solvent chosen (see 8.5.1.1).

Remove the plate from the tank, allow the solvents to evaporate from the plate in the dark and then examine under UV light, placing the plate 10 cm from the lamp (6.8). The spots of aflatoxin B<sub>1</sub> show a blue fluorescence.

### 8.5.2 Method B — Two-dimensional thin-layer chromatography

#### 8.5.2.1 Application of the solutions (see figure 1)

Trace two straight lines on a plate (6.7) parallel to two contiguous sides (50 and 60 mm from each side respectively), to establish the limit of migration of the solvent fronts. Apply the following solutions on the plate using capillary pipettes or microsyringes :

- at point A : 20 µl of the purified sample extract obtained in 8.4.2;

- at point B : 20 µl of the standard aflatoxin B<sub>1</sub> solution (5.14);
- at point C : 10 µl of the standard aflatoxin B<sub>1</sub> solution (5.14);
- at point D : 20 µl of the standard aflatoxin B<sub>1</sub> solution (5.14);
- at point E : 40 µl of the standard aflatoxin B<sub>1</sub> solution (5.14).

Dry in a slow stream of air or inert gas (5.12). The spots obtained shall have a diameter of about 5 mm.

#### 8.5.2.2 Development (see figure 1)

Develop the chromatogram in direction I, in the dark, using the developing solvent (5.6.3) (1 cm layer in a saturated tank) until the solvent front reaches the limit line. Remove the plate from the tank and allow to dry, in the dark, at ambient temperature for at least 15 min.

Then develop the chromatogram in direction II, in the dark, using the developing solvent (5.6.1) (1 cm layer in an unsaturated tank) until the solvent front reaches the limit line. Remove the plate from the tank and allow to dry, in the dark, at ambient temperature.

#### 8.5.2.3 Interpretation of the chromatogram (see figure 2)

Examine the chromatogram under UV light by placing the plate 10 cm from the lamp (6.8). Locate the position of the blue fluorescent spots B, C, D, and E of the aflatoxin B<sub>1</sub> from the standard solution and trace two imaginary lines passing through these spots and at right angles to the directions of development. The point P of intersection of these lines is the location at which to expect the aflatoxin B<sub>1</sub> spot originating from the test portion extract applied at point A (see figure 1). However, the actual location of the aflatoxin B<sub>1</sub> spot may be at a point Q at the intersection of two imaginary straight lines forming an angle of about 100° between them and passing through points B and C respectively.

#### 8.5.2.4 Supplementary chromatography

Trace on a new plate (6.7) two straight lines parallel to two contiguous sides, as indicated in figure 1, and apply, at point A, 20 µl of the purified test portion extract obtained in 8.4.2 and, superimposed on it, 20 µl of the standard aflatoxin B<sub>1</sub> solution (5.14). Develop as indicated in 8.5.2.2. Examine the chromatogram under UV light and check that

- the aflatoxin B<sub>1</sub> spots from the extract and the standard solution are superimposed;
- the fluorescence of this spot is more intense than that of the aflatoxin B<sub>1</sub> spot developed at point Q on the first plate.

## 8.6 Determination

Two methods of determination may be used, i.e. visual measurement or fluorodensitometric measurement.

### 8.6.1 Visual measurement

#### 8.6.1.1 Method A

Determine the quantity of aflatoxin B<sub>1</sub> in the extract by comparing the intensity of fluorescence of the extract spots with that of the standard solution spots. Interpolate if necessary.

The fluorescence obtained by the superimposition of the extract on the standard solution shall be more intense than that of the 10 µl of extract and shall be perceptible as only one spot. If the intensity of fluorescence given by the 10 µl of extract is greater than that of the 40 µl of standard solution, dilute the extract 10 or 100 times with the chloroform (5.1) or with the benzene/acetonitrile mixture (5.4) before repeating thin-layer chromatography.

#### 8.6.1.2 Method B

Determine the quantity of aflatoxin B<sub>1</sub> in the extract by comparing the intensity of the extract spot with that of spots C, D, and E from the standard solution. Interpolate if necessary.

If the intensity of fluorescence given by the 20 µl extract is greater than that of the 40 µl of standard solution, dilute the extract 10 or 100 times with the chloroform (5.1) or with the benzene/acetonitrile mixture (5.4) before repeating thin-layer chromatography.

### 8.6.2 Measurement by fluorodensitometry

Measure the intensity of fluorescence of the aflatoxin B<sub>1</sub> spots with the fluorodensitometer (6.10) at an excitation wavelength of 365 nm and an emission wavelength of 443 nm.

Determine, in the case of method A, the quantity of aflatoxin B<sub>1</sub> in the extract spots by comparison with the intensity of fluorescence of the spots from the standard solution, and, in the case of method B, the quantity of aflatoxin B<sub>1</sub> in the extract spot by comparison with the intensity of fluorescence of spots C, D and E from the standard solution.

### 8.7 Confirmation of the identity of aflatoxin B<sub>1</sub>

Confirm the identity of the aflatoxin B<sub>1</sub> in the extract by the presumptive test with sulphuric acid (see 8.7.1) and, if the result of this test is positive, by the actual confirmation test (8.7.2). If the result of the presumptive test with sulphuric acid (8.7.1) is negative, there is no need to proceed with the actual confirmation (8.7.2) since, in this case, no aflatoxin B<sub>1</sub> is present.

#### 8.7.1 Presumptive test with sulphuric acid

Spray the sulphuric acid (5.13) on to the chromatogram obtained in 8.5.1 or 8.5.2. The fluorescence of the aflatoxin B<sub>1</sub> spots shall turn from blue to yellow under UV light.

#### 8.7.2 Confirmation test: formation of aflatoxin B<sub>1</sub>-hemiacetal (aflatoxin B<sub>2a</sub>)

In the case of simple and only slightly pigmented feeds, use the one-dimensional thin-layer chromatographic method described

in 8.7.2.1. In the case of simple pigmented feeds, mixed feeds, or in cases of doubt, use the two-dimensional thin-layer chromatographic method described in 8.7.2.2.

#### 8.7.2.1 One-dimensional thin-layer chromatography

Trace a straight line on a plate (6.7) to divide it into two equal parts. Apply on each part, 20 mm from the lower edge and at intervals of 15 mm, the volumes indicated below of the standard aflatoxin B<sub>1</sub> solution and the extract:

- 25 µl of the standard aflatoxin B<sub>1</sub> solution (5.14);
- a volume of the extract obtained in 8.4.2 containing approximately 2,5 ng of aflatoxin B<sub>1</sub>;
- 25 µl of the standard aflatoxin B<sub>1</sub> solution (5.14) and, superimposed on it, a volume of the extract obtained in 8.4.2 containing approximately 2,5 ng of aflatoxin B<sub>1</sub>.

Apply to one of the two halves of the plate, superimposed on the spots previously applied, 1 to 2 µl of the trifluoroacetic acid (5.11). Dry in a stream of air at ambient temperature.

Develop the chromatogram, in the dark, using one of the developing solvents (5.6). The choice of the solvent shall be made beforehand. The solvent system shall ensure that the aflatoxin B<sub>1</sub>-hemiacetal (aflatoxin B<sub>2a</sub>) is clearly separated from interfering substances. The solvent front should travel about 120 mm.

Allow the solvents to evaporate in the dark, and then spray sulphuric acid (5.13) on to the part of the plate not previously treated with the trifluoroacetic acid. Examine the plate under UV light.

The identity of aflatoxin B<sub>1</sub> is confirmed if

- a) the R<sub>f</sub> value of the aflatoxin B<sub>1</sub> derivative originating from the extract corresponds with that from the standard solution;
- b) the aflatoxin B<sub>1</sub> derivative originating from the standard solution, superimposed on the extract, has a fluorescence more intense than the aflatoxin B<sub>1</sub> derivative originating from the extract.

Since fluorescent spots from the extract, having the same R<sub>f</sub> value as the aflatoxin B<sub>1</sub>-hemiacetal might lead to a false positive interpretation of the chromatogram, their presence shall be checked on the part of the plate treated with sulphuric acid.

In cases of doubt, confirmation by two-dimensional thin-layer chromatography (8.7.2.2) shall be used.

#### 8.7.2.2 Two-dimensional thin-layer chromatography (see figure 3)

##### 8.7.2.2.1 Application of the solutions

Trace two straight lines on a plate (6.7), parallel to two contiguous sides (60 mm from each side), to establish the limit of

migration of the solvent fronts. Apply the following solutions on the plate using capillary pipettes or microsyringes :

- at point A : a volume of purified extract from the sample, obtained in 8.4.2, containing about 2,5 ng of aflatoxin B<sub>1</sub>, and a drop (1 to 2 μl) of the trifluoroacetic acid (5.11);
- at points B and C : 25 μl of the standard aflatoxin B<sub>1</sub> solution (5.14), and a drop of the trifluoroacetic acid (5.11).

Dry in a stream of air at ambient temperature.

#### 8.7.2.2.2 Development

Develop the chromatogram in direction I, in the dark, using the developing solvent (5.6.2) (1 cm layer in an unsaturated tank) until the solvent front reaches the limit line. Remove the plate from the tank and allow to dry, in the dark, at ambient temperature, for 5 min.

Then develop the chromatogram in direction II, in the dark, using the developing solvent (5.6.1) (1 cm layer in an unsaturated tank) until the solvent front reaches the limit line. Remove the plate from the tank and allow to dry at ambient temperature.

#### 8.7.2.2.3 Interpretation of the chromatogram

Examine the chromatogram under UV light from the lamp (6.8) and check for the following features :

- a) appearance of a blue fluorescent spot of aflatoxin B<sub>1</sub>-hemiacetal, and sometimes a weak blue fluorescent spot of aflatoxin B<sub>1</sub> which has not reacted with the trifluoroacetic acid, originating from the standard solution applied at point C (migration in direction I) and from the standard solution applied at point B (migration in direction II);
- b) appearance of spots similar to those described in a), originating from the sample extract applied at point A. The position of these spots is defined by those originating from the standard solution applied at points B and C. The intensities of fluorescence of the aflatoxin B<sub>1</sub>-hemiacetal spots originating from the extract and from the standard solution applied at points B and C should be comparable.

### 8.8 Number of determinations

Carry out two determinations on the same test sample.

## 9 Expression of results

### 9.1 Method of calculation and formula

#### 9.1.1 Visual measurements

The aflatoxin B<sub>1</sub> content, expressed in micrograms per kilogram of sample, is equal to

$$\frac{C \times V_1 \times V_3}{m \times V_2}$$

where

*C* is the concentration, in micrograms of aflatoxin B<sub>1</sub> per millilitre of the standard solution (5.14) (approximately 0,1 μg/ml);

*m* is the mass, in grams, of the test portion corresponding to the volume of extract subjected to column clean-up (10,0 g);

*V*<sub>1</sub> is the final volume of the extract, in microlitres, taking into account any dilution that was necessary;

*V*<sub>2</sub> and *V*<sub>3</sub> are, respectively, the volumes, in microlitres, of the extract and of the standard aflatoxin B<sub>1</sub> solution (5.14), applied on the plate, having similar intensities of fluorescence.

### 9.1.2 Fluorodensitometric measurements

The aflatoxin B<sub>1</sub> content, expressed in micrograms per kilogram of sample, is equal to

$$\frac{m_1 \times V_1}{m \times V_2}$$

where

*m* is the mass, in grams, of the test portion corresponding to the volume of extract subjected to column clean-up (10,0 g);

*m*<sub>1</sub> is the mass, in nanograms, of aflatoxin B<sub>1</sub> in the extract spot (proportional to the volume *V*<sub>2</sub>), deduced from the measurements;

*V*<sub>1</sub> is the final volume of the extract, in microlitres, taking into account any dilution that was necessary;

*V*<sub>2</sub> is the volume, in microlitres, of extract applied on the plate (10 or 20 μl).

## 9.2 Precision

Three round-robin trials, two of which were carried out at the international level (Nos. 1 and 2), on mixed feeding stuffs (method B) gave the results indicated in the table.

The 11 laboratories participating in trial 2 also analysed the sample by method A, its composition being suitable, and obtained results very similar to those when using method B, by visual or fluorodensitometric measurement.

## 10 Test report

The test report shall show the method used (A or B), the method of determination (visual or fluorodensitometric measurement) and the result obtained. It shall also mention all operating conditions not specified in this International Standard, or regarded as optional, as well as any circumstances that may have influenced the results.

The test report shall include all the information necessary for the complete identification of the sample.