

ISO

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION

ISO RECOMMENDATION R 660

CRUDE VEGETABLE OILS AND FATS

DETERMINATION OF ACIDITY

1st EDITION

February 1968

COPYRIGHT RESERVED

The copyright of ISO Recommendations and ISO Standards belongs to ISO Member Bodies. Reproduction of these documents, in any country, may be authorized therefore only by the national standards organization of that country, being a member of ISO.

For each individual country the only valid standard is the national standard of that country.

Printed in Switzerland

Also issued in French and Russian. Copies to be obtained through the national standards organizations.

STANDARDSISO.COM : Click to view the full PDF of ISO/R 660:1968

BRIEF HISTORY

The ISO Recommendation R 660, *Crude vegetable oils and fats – Determination of acidity*, was drawn up by Technical Committee ISO/TC 34, *Agricultural food products*, the Secretariat of which is held by the Magyar Szabványügyi Hivatal (MSZH).

Work on this question by the Technical Committee began in 1961 and led, in 1963, to the adoption of a Draft ISO Recommendation.

In March 1966, this Draft ISO Recommendation (No. 901) was circulated to all the ISO Member Bodies for enquiry. It was approved, subject to a few modifications of an editorial nature, by the following Member Bodies :

Argentina	Hungary	Romania
Australia	India	South Africa,
Belgium	Iran	Rep. of
Bulgaria	Ireland	Turkey
Chile	Israel	U.A.R.
Colombia	Italy	United Kingdom
Czechoslovakia	Netherlands	U.S.S.R
Finland	New Zealand	Yugoslavia
France	Norway	
Germany	Poland	

No Member Body opposed the approval of the Draft.

The Draft ISO Recommendation was then submitted by correspondence to the ISO Council which decided, in February 1968, to accept it as an ISO RECOMMENDATION.

STANDARDSISO.COM : Click to view the full PDF of ISO/R 660:1968

CRUDE VEGETABLE OILS AND FATS

DETERMINATION OF ACIDITY

1. SCOPE

This ISO Recommendation describes a method for the determination of the free fatty acid content of crude vegetable oils and fats. This can be expressed as an acid value or as conventional acidity.

2. DEFINITIONS

- 2.1 *Acid value.* Number of milligrammes of potassium hydroxide required to neutralize the free fatty acids in 1 g of the oil or fat.
- 2.2 *Acidity.* Conventional expression of the percentage of free fatty acids (see clauses 7.2 and 7.3).

3. PRINCIPLE

Solution of a known quantity of the oil or fat to be analysed in a mixture of ethanol and diethyl-ether, followed by titration of the free fatty acids present with an ethanolic solution of potassium hydroxide.

4. REAGENTS

- 4.1 *Mixture 1 + 1 v/v ethanol 95 % v/v and diethyl ether.* Neutralize exactly, just before use, by means of the ethanolic potassium hydroxide solution (4.2), in the presence of 0.3 ml of indicator (4.3) per 100 ml of this mixture (phenolphthalein or alkali blue 6 B, as the case may be (see section 6).
- 4.2 *Potassium hydroxide,* approximately 0.1 N or, if necessary, approximately 0.5 N solution in ethanol 95 % v/v. The exact concentration should be known, and checked immediately before use. Use a solution prepared at least five days previously and decanted into a bottle of brown glass, provided with a rubber stopper. The solution should be colourless or straw yellow.

NOTE. — It is recommended that the ethanol should be purified by adding 5 to 10 g of potassium hydroxide to 1 litre of the ethanol, boiling for one hour under reflux and finally distilling over.

- 4.3 *Indicator.* Phenolphthalein, 10 g per litre of ethanol 95 % v/v or, if necessary, alkali blue 6 B, 20 g per litre of ethanol 95 % v/v.

5. APPARATUS

- 5.1 *Conical flask, 250 ml.*
- 5.2 *Burette graduated in 0.1 ml, complying with the specification for class A of ISO Recommendation R 385, Burettes.*
- 5.3 *Analytical balance.*

6. PROCEDURE

6.1 Preparation of sample

Prepare the contract sample as described in ISO Recommendation R 661, *Crude vegetable oils and fats – Preparation of contract sample for analysis.*

6.2 Test portion

Weigh into the 250 ml conical flask, to the nearest 0.01 g, 5 to 10 g of the oil or fat, according to the acidity expected.

6.3 Determination

Dissolve the test portion in about 150 ml of the 1 + 1 mixture of ethanol and diethyl ether (4.1), previously neutralized.

If the solution obtained is not perfectly clear, add a further quantity of the mixture (4.1).

Titrate, with shaking, with the 0.1 N ethanolic solution of potassium hydroxide (4.2) to the end point of the indicator (pink colour of phenolphthalein persisting for at least 10 seconds). If the quantity of 0.1 N potassium hydroxide solution required exceeds 20 ml, a 0.5 N solution should be used (see clauses 8.1, 8.2, 8.3).

Carry out two determinations on the same prepared sample.

7. EXPRESSION OF RESULTS

- 7.1 It is recommended that the result of the analysis should be expressed as an acid value (see clause 2.1). Since 1 ml of N potassium hydroxide solution corresponds to 56.1 mg of potassium hydroxide,

$$\text{acid value} = \frac{V \times 56.1}{M}$$

where

V is the volume, expressed as millilitres of N solution, of ethanolic potassium hydroxide solution used,

M is the mass, in grammes, of the test portion.

Take as a result the arithmetic means of two determinations.