

ISO

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION

ISO RECOMMENDATION R 979

SODIUM HYDROXIDE FOR INDUSTRIAL USE

DETERMINATION OF ALKALINITY

VOLUMETRIC METHOD

1st EDITION

February 1969

COPYRIGHT RESERVED

The copyright of ISO Recommendations and ISO Standards belongs to ISO Member Bodies. Reproduction of these documents, in any country, may be authorized therefore only by the national standards organization of that country, being a member of ISO.

For each individual country the only valid standard is the national standard of that country.

Printed in Switzerland

Also issued in French and Russian. Copies to be obtained through the national standards organizations.

STANDARDSISO.COM : Click to view the full PDF of ISO/R 979:1969

BRIEF HISTORY

The ISO Recommendation R 979, *Sodium hydroxide for industrial use – Determination of alkalinity – Volumetric method*, was drawn up by Technical Committee ISO/TC 47, *Chemistry*, the Secretariat of which is held by the Ente Nazionale Italiano di Unificazione (UNI).

Work on this question led, in 1966, to the adoption of a Draft ISO Recommendation.

In December 1966, this Draft ISO Recommendation (No. 1088) was circulated to all the ISO Member Bodies for enquiry. It was approved, subject to a few modifications of an editorial nature, by the following Member Bodies :

Austria	Ireland	South Africa, Rep. of
Belgium	Israel	Spain
Chile	Italy	Switzerland
Cuba	Japan	Thailand
Czechoslovakia	Korea, Dem. P. Rep. of	Turkey
France	Netherlands	U.A.R.
Germany	New Zealand	United Kingdom
Hungary	Poland	U.S.S.R.
India	Portugal	Yugoslavia
Iran	Romania	

One Member Body opposed the approval of the Draft :

U.S.A.

The Draft ISO Recommendation was then submitted by correspondence to the ISO Council, which decided, in February 1969, to accept it as an ISO RECOMMENDATION.

STANDARDSISO.COM : Click to view the full PDF of ISO/R 979:1969

SODIUM HYDROXIDE FOR INDUSTRIAL USE

DETERMINATION OF ALKALINITY

VOLUMETRIC METHOD

INTRODUCTION

Two expressions are commonly used for expressing the alkalinity of the product :

- *Equivalent* NaOH (eq. NaOH), corresponding to the total alkalinity of the product expressed as NaOH;
- *Caustic* NaOH (c. NaOH), corresponding to the total alkalinity of the product less that due to sodium carbonate.

PART I

DETERMINATION OF TOTAL ALKALINITY

1. SCOPE

Part I of this ISO Recommendation describes a volumetric method for the determination of the total alkalinity of sodium hydroxide for industrial use, expressed as equivalent NaOH (eq. NaOH).

2. PRINCIPLE

Neutralization of an aliquot of the sample solution by means of a hydrochloric acid standard volumetric solution in the presence of methyl orange as indicator, or any other indicator having the same equivalence point.

3. REAGENTS

Distilled water or water of equivalent purity should be used in the test.

3.1 *Hydrochloric acid*, N standard volumetric solution (see Note in section 6).

3.2 *Methyl orange*, 0.5 g/l solution.

Dissolve 0.05 g of methyl orange in water and dilute to 100 ml.

4. APPARATUS

Ordinary laboratory apparatus.

5. PROCEDURE

5.1 Test portion

Transfer 20.0 ml of sample solution A* to a 500 ml conical flask.

5.2 Titration

Add approximately 80 ml of water and 5 drops of the methyl orange solution (3.2) to the conical flask containing the test portion (5.1) and titrate with the hydrochloric acid standard volumetric solution (3.1) until the colour of the indicator changes from yellow to orange.

6. EXPRESSION OF RESULTS

The total alkalinity, expressed as equivalent sodium hydroxide (eq. NaOH), is given, as a percentage by mass, by the following formula :

$$V \times A \times \frac{500}{20} \times \frac{100}{E} = 99.9925 \times \frac{V}{E}$$

where

- V* is the volume, in millilitres, of the hydrochloric acid standard volumetric solution (3.1) used for the titration;
- A* is the mass, in grammes, of NaOH corresponding to 1 ml of N hydrochloric acid standard volumetric solution (theoretical value 1 ml \cong 0.039 997 g of NaOH);
- E* is the mass, in grammes, of the test portion used for the preparation of the sample solution A*.

NOTE. — If the hydrochloric acid standard volumetric solution (3.1) is not of exactly the strength indicated in the list of reagents, a suitable correction factor should be employed in calculating the results.

7. TEST REPORT

Give the following particulars :

- (a) the reference of the method used;
- (b) the results and the method of expression used;
- (c) any unusual features noted during the determination;
- (d) any operation not included in this ISO Recommendation or regarded as optional.

* See ISO Recommendation R 978, *Sodium hydroxide for industrial use — Preparation of sample solution*, clause 5.2.