

---

---

**Food products — Guidance on how to  
express vitamins and their vitamers  
content**

*Produits alimentaires — Lignes directrices pour exprimer les teneurs  
en vitamines et en leurs vitamères*

STANDARDSISO.COM : Click to view the full PDF of ISO/TR 23304:2021



STANDARDSISO.COM : Click to view the full PDF of ISO/TR 23304:2021



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2021

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
CP 401 • Ch. de Blandonnet 8  
CH-1214 Vernier, Geneva  
Phone: +41 22 749 01 11  
Email: [copyright@iso.org](mailto:copyright@iso.org)  
Website: [www.iso.org](http://www.iso.org)

Published in Switzerland

# Contents

	Page
Foreword.....	iv
Introduction.....	v
<b>1 Scope.....</b>	<b>1</b>
<b>2 Normative references.....</b>	<b>1</b>
<b>3 Terms and definitions.....</b>	<b>1</b>
<b>4 Guidelines per vitamins.....</b>	<b>1</b>
4.1 General.....	1
4.2 Vitamin E.....	2
4.3 Vitamin A.....	5
4.4 Vitamin D.....	7
4.5 Vitamin K.....	8
4.6 Thiamine (vitamin B <sub>1</sub> ).....	10
4.7 Riboflavin (vitamin B <sub>2</sub> ).....	11
4.8 Niacin (vitamin B <sub>3</sub> or vitamin PP).....	12
4.9 Vitamin B <sub>6</sub> .....	12
4.10 Biotin (vitamin H or vitamin B <sub>8</sub> ).....	13
4.11 Folates (vitamin B <sub>9</sub> ).....	14
4.12 Vitamin B <sub>12</sub> .....	15
4.13 Pantothenic acid (vitamin B <sub>5</sub> ).....	16
4.14 Vitamin C.....	17
<b>5 US FDA unit conversions.....</b>	<b>19</b>
<b>Bibliography.....</b>	<b>20</b>

STANDARDSISO.COM : Click to view the full PDF of ISO/TR 23304:2021

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see [www.iso.org/directives](http://www.iso.org/directives)).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see [www.iso.org/patents](http://www.iso.org/patents)).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see [www.iso.org/iso/foreword.html](http://www.iso.org/iso/foreword.html).

This document was prepared by Technical Committee ISO/TC 34, *Food products*.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at [www.iso.org/members.html](http://www.iso.org/members.html).

## Introduction

Vitamins can be naturally found in foods in different molecular forms. In more elaborated food products, vitamins can be used for fortification by adding several molecular forms with different levels of vitamin activity. There are regulations governing the addition of vitamins in food products. The authorized compounds for fortification depend on the type of food. Regulation deals with, for instance baby food or food supplements. The main problem is that the vitamin activities of the authorized compounds are not clearly described.

At the same time, it is not mandatory to list the chemical name of the compound used for food fortification purposes according to food labelling regulations. For example, vitamin E can be written in the list of ingredients without knowing if it is D-alpha tocopherol or D,L-alpha tocopherol, even though each molecular form presents different vitamin E activity.

ISO and CEN analytical standards express results in mass units related to the vitamin standard used for quantification. As expression in specific units of vitamin activity can be linked to regional/national regulatory requirements, analytical methods do not give guidance for this conversion. This document proposes ways to express the vitamin content in order to facilitate harmonization between different laboratories and also to reduce misunderstanding of the results expressed in vitamin content and in some cases in the vitamin activity of vitamins.

STANDARDSISO.COM : Click to view the full PDF of ISO/TR 23304:2021

[STANDARDSISO.COM](https://standardsiso.com) : Click to view the full PDF of ISO/TR 23304:2021

# Food products — Guidance on how to express vitamins and their vitamers content

## 1 Scope

This document provides guidelines on:

- how to express vitamin quantity,
- the expression of different molecular forms in appropriate units,
- and in some cases, vitamin activity, according to vitamers present or used in food products, in addition to the quantitative content determination obtained from ISO and CEN analytical standards.

It provides information to be used as a basis for discussion between stakeholders and food control laboratories. It is not intended to be prescriptive or exhaustive.

## 2 Normative references

There are no normative references in this document.

## 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <http://www.electropedia.org/>

### 3.1

#### vitamer

any of a number of chemical compounds of a particular vitamin, generally having a similar molecular structure, each of which shows vitamin activity in a vitamin-deficient biological system

Note 1 to entry: See Reference [1].

## 4 Guidelines per vitamins

### 4.1 General

This document begins by addressing vitamin E, as this vitamin is considered to be the most complex in terms of variation of possible units of measurement. Vitamin E is followed by the other fat-soluble vitamins and water-soluble vitamins respectively.

To include the diversity of available conversion factors, regulations in different countries/regions are evaluated, including the European Union (EU), Regulations of the United States Food and Drug Administration (USFDA), United States of America (USA), National Standard (GB) of China, Food Standards Australia/New Zealand (FSANZ). In addition, standards of global organizations (CODEX<sup>[2]</sup>) are included.

## 4.2 Vitamin E

Vitamin E is a fat-soluble vitamin. Vitamin E is mainly found in the fat of the different food products as tocopherols and tocotrienols esters.

Food regulations of many countries authorize the fortification of different kind of food products with molecules having a vitamin E activity. The names of the authorized molecular forms to add are generally linked to the type of food product. [Table 1](#) gives an example of vitamin E compounds permitted for use for food fortification according to EU regulations, depending on the product type.

**Table 1 — Example of authorized vitamin E compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	D-alpha-tocopherol	430,7	Regulation (EU) n° 609/2013 <sup>[3]</sup>
	DL-alpha-tocopherol	430,7	
	D-alpha-tocopheryl acetate	472,8	
	DL-alpha-tocopheryl acetate	472,8	
	D-alpha-tocopheryl polyethylene glycol 1000 succinate		
Supplemented food	D-alpha-tocopherol	430,7	Regulation (EC) n°1925/2006 <sup>[4]</sup>
	DL-alpha-tocopherol	430,7	
	D-alpha-tocopheryl acetate	472,8	
	DL-alpha-tocopheryl acetate	472,8	
	D-alpha-tocopheryl acid succinate	530,8	
Food supplements	D-alpha-tocopherol	430,7	Directive n°2002/46/EC <sup>[5]</sup>
	DL-alpha-tocopherol	430,7	
	D-alpha-tocopheryl acetate	472,8	
	DL-alpha-tocopheryl acetate	472,8	
	D-alpha-tocopheryl acid succinate	530,8	
	Mix of tocopherols and tocotrienols		

There is no global convention on how the amount of vitamin E should be expressed for labelling purposes. In [Table 2](#) an overview of units of measurement for vitamin E is given according to several national/regional regulations and a CODEX standard.

**Table 2 — Definition of vitamin E for labelling purposes in several standards/regulations**

US (until 2019-12-31)	US (as from 2020-01-01)	EU	CODEX	China/Australia/New Zealand
IU <sup>a</sup>	mg alpha-tocopherol	mg alpha - TE <sup>b</sup>	mg alpha - TE	mg alpha - TE
<sup>a</sup> International Unit, a measure of biological activity, different for each substance. <sup>b</sup> Tocopherol equivalent.				

Table 2 (continued)

US (until 2019-12-31)	US (as from 2020-01-01)	EU	CODEX	China/Australia/New Zealand
1 IU = 0,67 mg for D-alpha tocopherol (natural)	1 mg vitamin E (as alpha-tocopherol) label claim = 1 mg of natural alpha-tocopherol (D-alpha-tocopherol)	1 mg alpha - TE = 1 mg D-alpha tocopherol	1 mg alpha - TE = 1 mg D-alpha tocopherol	1 mg alpha - TE = 1 mg D-alpha tocopherol
1 IU = 0,9 mg for D, L-alpha tocopherol (synthetic)	1 mg vitamin E (as alpha-tocopherol) label claim = 2 mg of synthetic alpha-tocopherol (D,L-alpha-tocopherol)			
<p><sup>a</sup> International Unit, a measure of biological activity, different for each substance.</p> <p><sup>b</sup> Tocopherol equivalent.</p>				

The vitamin E unit of measurement is generally defined as 1 mg of D-alpha-tocopherol (natural form). There is currently no standard or official analytical method able to quantify this natural form specifically.

There are two main analytical approaches for quantifying vitamin E:

- 1) Saponification of tocopheryl esters to liberate them as free tocopherols. Solvent extraction and separation of the individual tocopherols by liquid chromatography. Quantification by fluorescence detection.

An example is EN 12822[6], which includes a saponification and an extraction step. It enables the separation by a normal phase HPLC, and the quantification of the free alpha, beta, gamma and delta tocopherols using fluorescence detection. Tocotrienols are also separated but not quantified. Tocotrienols are not taken into account for the expression of the vitamin result. The D-tocopherols are not distinguished from the D,L-tocopherols.

- 2) Direct solvent extraction and separation of different esters and free tocopherols by liquid chromatography. Quantification by fluorescence detection.

An example is ISO 20633, where the hydrophilic coating of fat micelles is hydrolysed by an enzyme. The hydrophobic contents of the micelles are then extracted into iso-octane. The extract is analysed by normal phase HPLC. Alpha-tocopherol and alpha-tocopheryl acetate are quantified using fluorescence detection.

By taking into account the molecular mass and the activity ratio of the D and the D,L the following values can be found<sup>[8]</sup>:

- 1 mg of vitamin E = 1 mg of D-alpha-tocopherol = 1 mg alpha-TE;
- 1 mg of vitamin E = 1,35 mg of D,L-alpha-tocopherol;
- 1 mg of vitamin E = 1,1 mg of D-alpha-tocopheryl acetate;
- 1 mg of vitamin E = 1,49 mg of D,L-alpha-tocopheryl acetate;
- 1 mg of vitamin E = 1,23 mg of D-alpha-tocopheryl acid succinate.

As mentioned above, there is no specific detection at the quantification step for the D-alpha tocopherol (natural form) and the D,L-alpha tocopherol (synthetic form) respectively. So, when a sample is analysed, and there is no specific information about the added form (D or D,L) on the label of the product, there

can be a misunderstanding about the true value of the vitamin E content. The difference can be a factor of 1,35 or higher.

Regarding the nutritional aspect, there is no general agreement about the activities of the other forms of tocopherols and tocotrienols.

When in the presence of certain amounts of beta and/or gamma and/or delta tocopherols, it is often underlined that the relative vitamin E activities should be included in the final result. The more common coefficients<sup>[8]</sup> are:

- 1 mg of D-beta-tocopherol = 0,5 mg of vitamin E;
- 1 mg of D-gamma-tocopherol = 0,1 mg of vitamin E;
- 1 mg of D-delta-tocopherol = 0,03 mg of vitamin E.

Tocotrienols are sometimes also taken into account by certain guidelines, such as the FAO/INFOODS guideline<sup>[9]</sup> as in the following example:

Vitamin E (mg/100 g) = alpha-tocopherol (mg/100 g) + 0,4 beta-tocopherol (mg/100 g) + 0,1 gamma-tocopherol (mg/100 g) + 0,01 delta-tocopherol (mg/100 g) + 0,3 alpha-tocotrienol (mg/100 g) + 0,05 beta-tocotrienol (mg/100 g) + 0,01 gamma-tocotrienol (mg/100 g)

Despite this fact, in most of the cases, only the alpha-tocopherol is considered.

At the session of the Codex Committee on Nutrition and Foods for Special Dietary Uses (40<sup>th</sup> Session, 5<sup>th</sup> - 9<sup>th</sup> December 2016, Hamburg, Germany), it was proposed to take only the alpha-tocopherol into account for labelling purposes<sup>[10]</sup>. In this case, the expression of vitamin E is very simple:

vitamin E = 1 mg of alpha-D-tocopherol = 1 mg RRR-alpha-tocopherol.

Taking into account most of these parameters, the vitamin E content can be calculated as follows:

- a) If only D-alpha tocopherol (natural) is present in the product, the vitamin E content (expressed in D-alpha tocopherol equivalent) can be calculated as vitamin E (mg/100 g) = alpha-D-tocopherol

or in some cases, e.g. for vegetable oils:

vitamin E (mg/100 g) = alpha-tocopherol + 0,5 beta-tocopherol + 0,1 gamma-tocopherol + 0,03 delta-tocopherol.

- b) If D,L alpha tocopherol (synthetic) is present in the product, the alpha-tocopherol result is divided by 1,35 and the vitamin E content (expressed in D-alpha tocopherol equivalent) can be calculated as vitamin E (mg/100 g) = 0,74 alpha-tocopherol (mg/100 g)

or in some cases:

vitamin E (mg/100 g) = 0,74 alpha-tocopherol (mg/100 g) + 0,5 beta-tocopherol (mg/100 g) + 0,1 gamma-tocopherol (mg/100 g) + 0,03 delta-tocopherol (mg/100 g).

- c) If the vitamin E molecular form is not known, it is recommended that the two former possibilities be studied. Follow-up action with the producer of the product can be necessary before a conclusion can be drawn.

- d) Some methods do not involve a saponification step, like in ISO 20633. In these cases, the calculation is adapted as follows:

vitamin E (mg/100 g) = D-alpha tocopherol (mg/100 g) + 0,909 D-alpha tocopherol acetate (mg/100 g).

In addition, it is important to consider that apart from adding vitamin E for food fortification purposes, it is common practice and permitted to add tocopherols as food additives (antioxidants) to food products, including:

- E 306: rich tocopherols extracts;
- E 307: alpha tocopherol;
- E 308: gamma tocopherol;
- E 309: delta tocopherol.

At the analysis step, this can lead to a difference between the vitamin E quantified value and the vitamin E labelled value because no test is capable of identifying the part of vitamin E added for nutritional purposes from the part added for antioxidation. In any case, the labelled vitamin E value should not include the tocopherol forms added as antioxidants.

### 4.3 Vitamin A

Vitamin A is a fat-soluble vitamin. Vitamin A is mainly found in the fat of the different food products as retinol esters.

The food regulations of many countries authorize the fortification of different kinds of food products with molecules having a vitamin A activity. The names of the authorized molecular forms to add in are generally linked to the type of food product. [Table 3](#) gives an example of vitamin A compounds permitted for use for food fortification according to EU regulations, depending on the product type.

**Table 3 — Example of authorized vitamin A compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	Retinol	286,5	Regulation (EU) n° 609/2013
	Retinyl acetate	328,5	
	Retinyl palmitate	524,9	
	Beta carotene	536,9	
Supplemented food	Retinol	286,5	Regulation (EC) n° 1925/2006
	Retinyl acetate	328,5	
	Retinyl palmitate	524,9	
	Beta carotene	536,9	
Food supplements	Retinol	286,5	Directive n°2002/46/EC
	Retinyl acetate	328,5	
	Retinyl palmitate	524,9	
	Beta carotene	536,9	

There is no global convention on how the amount of vitamin A should be expressed for labelling purposes. In [Table 4](#) an overview of units of measurement for vitamin A is given according to several national/regional regulations and a CODEX standard.

Table 4 — Definition of vitamin A for labelling purposes in several standards/regulations

US (until 2019-12-31)	US (as from 2020-01-01)	EU	CODEX	China/Australia/New Zealand
IU	$\mu\text{g RAE}^a$	$\mu\text{g RE}$	$\mu\text{g RE}$	$\mu\text{g RE}$
1 IU = 0,3 $\mu\text{g}$ retinol	1 $\mu\text{g RAE}$ = 1 $\mu\text{g}$ retinol	1 $\mu\text{g RE}$ = 1 $\mu\text{g}$ retinol	1 $\mu\text{g RE}$ = 1 $\mu\text{g}$ retinol 1 $\mu\text{g}$ retinol = 3,33 IU	1 $\mu\text{g RE}$ = 1 $\mu\text{g}$ retinol 1 $\mu\text{g}$ retinol = 3,33 IU
1 IU = 0,6 $\mu\text{g}$ of beta-carotene	1 $\mu\text{g RAE}$ = 2 $\mu\text{g}$ supplemental beta-carotene	6 $\mu\text{g}$ beta-carotene = 1 $\mu\text{g RE}$	Carotenoids excluded	Carotenoids excluded
	1 $\mu\text{g RAE}$ = 12 $\mu\text{g}$ beta-carotene			
	1 $\mu\text{g RAE}$ = 24 $\mu\text{g}$ alpha-carotene			
	1 $\mu\text{g RAE}$ = 24 $\mu\text{g}$ beta-cryptoxanthin			
<sup>a</sup> Retinol activity equivalent. <sup>b</sup> Retinol equivalent.				

In general, the unit of measurement for vitamin A is  $\mu\text{g RE}$  and defined as 1  $\mu\text{g}$  of retinol. The natural form is the trans retinol.

There are two main analytical approaches for quantifying vitamin A:

- 1) Saponification of retinol esters to liberate them as free retinol. Solvent extraction and separation of the cis and trans retinol by liquid chromatography. Quantification by UV or fluorescence detection.

An example is EN 12823-1<sup>[11]</sup>, which includes a saponification step and enables the separation and the quantification of all E-retinol (trans-retinol) and 13-Z-retinol (13-cis-retinol). EN 12823-2<sup>[12]</sup> enables the quantification of beta-carotene.

- 2) Direct solvent extraction and separation of different esters of retinol and retinol by liquid chromatography. Quantification by UV or fluorimetric detection.

An example is ISO 20633, where the hydrophilic coating of fat micelles is hydrolysed by an enzyme. The hydrophobic contents of the micelles are then extracted into iso-octane. The extract is analysed by normal phase HPLC. Vitamin A palmitate (cis and trans) and vitamin A acetate (cis and trans) are quantified using UV detection.

By taking into account the molecular mass, the following values for retinyl esters can be found:

- 1  $\mu\text{g}$  of all trans retinyl acetate = 0,872  $\mu\text{g RE}$  (vitamin A);
- 1  $\mu\text{g}$  of all trans retinyl palmitate = 0,546  $\mu\text{g RE}$  (vitamin A).

In some regulations, beta-carotene can be included as vitamin A. In that case the beta-carotene content value should be transformed in RE. The correction factor is in most of the cases: 1  $\mu\text{g}$  of retinol = 6  $\mu\text{g}$  of beta-carotene.

In the literature, different factors for alpha-carotene and/or some other carotenoids can be taken into account like in the Souci-Fachmann-Kraut food composition and nutrition tables<sup>[13]</sup>:

vitamin A = retinol + 1/6 beta-carotene + 1/12 (other vitamin A active carotenoids).

Other vitamin A active carotenoids include alpha-carotene, gamma-carotene, cryptoxanthin and mutatochrome.

These carotenoids are usually not taken into account because of their low activity in comparison to beta-carotene and because of the need for additional assays.

To take into account most of these parameters, the vitamin A content can be calculated as:

$$\text{vitamin A } (\mu\text{g}/100 \text{ g}) = \text{retinol } (\mu\text{g}/100 \text{ g}) [\text{all E-retinol (trans-retinol) + 13-Z-retinol (13-cis-retinol)}]$$

or in some cases, depending on the regulation:

$$\text{vitamin A } (\mu\text{g}/100 \text{ g}) = \text{retinol } (\mu\text{g}/100 \text{ g}) + 0,167 \text{ beta-carotene } (\mu\text{g}/100 \text{ g}).$$

Some methods do not involve a saponification step, like in ISO 20633. In these cases, the calculation is adapted as follows:

$$\text{vitamin A as retinol } (\mu\text{g}/100 \text{ g}) = 0,55 \times \text{retinyl palmitate } (\mu\text{g}/100\text{g}) + 0,87 \times \text{retinyl acetate } (\mu\text{g}/100 \text{ g}).$$

#### 4.4 Vitamin D

Vitamin D is a fat-soluble vitamin. Vitamin D is found in the fat of many food products. Vitamin D<sub>2</sub> is found in vegetables and vitamin D<sub>3</sub> is found in food from animal origin (for example meat, fish and milk).

Food regulations of many countries authorize the fortification of different kinds of food products with molecules having a vitamin D activity. The names of the authorized molecular forms to add in are generally linked to the type of food product. [Table 5](#) gives an example of vitamin D forms permitted for use for food fortification according to EU regulations, depending on the product type.

**Table 5 — Example of authorized vitamin D compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	Vitamin D <sub>2</sub>	396,7	Regulation (EU) n° 609/2013
	Vitamin D <sub>3</sub>	384,6	
Supplemented food	Vitamin D <sub>2</sub>	396,7	Regulation (EC) n° 1925/2006
	Vitamin D <sub>3</sub>	384,6	
Food supplements	Vitamin D <sub>2</sub>	396,7	Directive n°2002/46/EC
	Vitamin D <sub>3</sub>	384,6	

There are several vitamers of vitamin D. The two major forms are vitamin D<sub>2</sub> (ergocalciferol) and vitamin D<sub>3</sub> (cholecalciferol). Vitamin D without a subscript refers to either vitamin D<sub>2</sub> or vitamin D<sub>3</sub> or both and is known as calciferol. There is global agreement about the units and the conversion factors to be used. In [Table 6](#) an overview of units of measurement for vitamin D is given according to several national/regional regulations and a CODEX standard.

**Table 6 — Definition of vitamin D for labelling purposes in several standards/regulations**

US (until 2019-12-31)	US (as from 2020-01-01)	EU	CODEX	China / Australia / New Zealand
IU	μg	μg	μg	μg
1 IU = 0,025 μg ergocalciferol or cholecalciferol	1 μg = 40 IU	1 μg vitamin D <sub>2</sub> or 1 μg vitamin D <sub>3</sub> (or 1 μg vitamin D <sub>2</sub> + vitamin D <sub>3</sub> )	1 μg vitamin D <sub>3</sub>	1 μg vitamin D <sub>2</sub> or 1 μg vitamin D <sub>3</sub> (or 1 μg vitamin D <sub>2</sub> +vitamin D <sub>3</sub> )
vitamin D <sub>2</sub> and/or vitamin D <sub>3</sub>	vitamin D <sub>2</sub> and/or vitamin D <sub>3</sub>	vitamin D <sub>2</sub> and/or vitamin D <sub>3</sub>	vitamin D <sub>3</sub>	vitamin D <sub>2</sub> and/or vitamin D <sub>3</sub>

In general, the unit of measurement for vitamin D is  $\mu\text{g}$  cholecalciferol or ergocalciferol.

It is considered that for vitamin D, 1  $\mu\text{g}$  of ergocalciferol (vitamin D<sub>2</sub>) is equivalent to 1  $\mu\text{g}$  cholecalciferol (vitamin D<sub>3</sub>) because both have the same vitamin D activity.

According to former US regulation, the International Unit (IU) of vitamin D is equal to 0,025  $\mu\text{g}$  of vitamin D<sub>2</sub> or vitamin D<sub>3</sub>. The IU can be still be found on labels.

There are two main analytical approaches for quantifying vitamin D:

- 1) Saponification of the matrix to liberate vitamin D<sub>2</sub> and vitamin D<sub>3</sub> as free forms. Solvent extraction. A first HPLC step to purify and collect the vitamin D<sub>2</sub> and vitamin D<sub>3</sub> fraction. Then separation of the vitamin D<sub>2</sub> and vitamin D<sub>3</sub> by liquid chromatography. Quantification by UV detection.

An example is EN 12821<sup>[14]</sup>, which includes a saponification step. A first HPLC step enables the purification and collection of vitamins D<sub>2</sub> and D<sub>3</sub>. Then vitamin D<sub>2</sub> and vitamin D<sub>3</sub> are separated by a second chromatography. Both are quantified by UV detection. The drawback of this method is that vitamin D<sub>2</sub> and vitamin D<sub>3</sub> cannot be quantified simultaneously because one is used as an internal standard to quantify the other.

- 2) Saponification of the matrix to liberate vitamin D<sub>2</sub> and vitamin D<sub>3</sub> as free forms. Derivatization of the vitamin D to a higher molecular mass. Then separation by HPLC and quantification by mass spectrometry. The advantage of this principle is that vitamin D<sub>2</sub> and vitamin D<sub>3</sub> can be quantified simultaneously.

An example is ISO 20636 where samples are saponified and vitamin D is extracted by isooctane. Vitamin D is derivatized and then quantified by mass spectrometry after a chromatographic step. Stable isotopes of vitamin D<sub>2</sub> and vitamin D<sub>3</sub> are used as internal standards.

To conclude, the vitamin D content can be calculated as:

$$\text{vitamin D } (\mu\text{g}/100 \text{ g}) = \text{ergocalciferol } (\mu\text{g}/100 \text{ g}) + \text{cholecalciferol } (\mu\text{g}/100 \text{ g}).$$

#### 4.5 Vitamin K

Vitamin K is a fat-soluble vitamin. Vitamin K can be found as vitamin K<sub>1</sub> in vegetables (spinach, lettuce, broccoli, Brussels sprouts) and as vitamin K<sub>1</sub> and vitamin K<sub>2</sub> in meat and liver.

Food regulations of many countries authorize the fortification of different kind of food products with molecules having a vitamin K activity. The names of the authorized molecular forms to add in are generally linked to the type of food product. [Table 7](#) gives an example of vitamin K compounds permitted for use for food fortification according to EU regulations depending the product type.

**Table 7 — Example of authorized vitamin K compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	Phylloquinone (phytomenadione - vitamin K <sub>1</sub> )	450,7	Regulation (EU) n° 609/2013
	Menaquinone (vitamin K <sub>2</sub> )		
	K <sub>2</sub> -MK <sub>4</sub>	444,6	
	K <sub>2</sub> -MK <sub>7</sub>	649,0	

Table 7 (continued)

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Supplemented food	Phylloquinone (phytomenadione – vitamin K <sub>1</sub> )	450,7	Regulation (EC) n° 1925/2006
	Menaquinone (vitamin K <sub>2</sub> )		
	K <sub>2</sub> -MK <sub>4</sub>	444,6	
	K <sub>2</sub> -MK <sub>7</sub>	649,0	
Food supplement products	Phylloquinone (phytomenadione – vitamin K <sub>1</sub> )	450,7	Directive n°2002/46/EC
	Menaquinone (vitamin K <sub>2</sub> )		
	K <sub>2</sub> -MK <sub>4</sub>	444,6	
	K <sub>2</sub> -MK <sub>7</sub>	649,0	

There are three main vitamers of vitamin K. The major form is vitamin K<sub>1</sub> (phylloquinone or phytomenadione or phytonadione). The other forms are vitamin K<sub>2</sub> (menaquinone) and vitamin K<sub>3</sub> (menadione).

In general, the unit of measurement for vitamin K is µg vitamin K<sub>1</sub> (phylloquinone) or vitamin K<sub>2</sub> (menaquinone).

Menaquinone (vitamin K<sub>2</sub>) can also be added to different kinds of food in different molecular forms (Table 7). The difference in molecular form is linked to the length of the isoprenoid chain. The number of isoprenoid groups gives the name of the menaquinone. With the 4 isoprenoid group the name is vitamin K<sub>2</sub>-MK<sub>4</sub> and with the 7 isoprenoid group the name is vitamin K<sub>2</sub>-MK<sub>7</sub>.

To conclude, the vitamin K content can be calculated as:

Vitamin K (µg/100 g) = vitamin K<sub>1</sub> (µg/100 g), or vitamin K<sub>2</sub> (µg/100 g).

It should be noted that a significant level of cis isomer can be found in synthetic vitamin K<sub>1</sub>. Cis-vitamin K<sub>1</sub> is considered less bioavailable. However, food regulations do not distinguish between cis- and trans-vitamin K<sub>1</sub>. From a food composition perspective, it can be more relevant to quantify trans-vitamin K<sub>1</sub>, whereas from a regulatory/food labelling perspective total vitamin K<sub>1</sub> should be considered.

Most of methods quantify total vitamin K<sub>1</sub>, but the recent IS ISO 21446 enables the quantification of the cis and the trans isomers separately.

In ISO 21446, vitamin K<sub>1</sub> is extracted from products with iso-octane after precipitation of proteins and release of lipids with methanol. Prepared samples are injected on a direct phase HPLC column where cis and trans vitamin K<sub>1</sub> are separated isocratically with an iso-octane-isopropanol mobile phase. The column eluent is mixed with a dilute ethanolic solution of zinc chloride, sodium acetate and acetic acid, and vitamin K<sub>1</sub> is reduced to a fluorescent derivative in a zinc reactor column. The resulting fluorescent compound is quantified with fluorimetric detection.

EN 14148<sup>[17]</sup>, for example, includes the destruction of lipids by an enzymatic treatment and a post-column reduction before a fluorimetric quantification. It enables the quantification of total vitamin K<sub>1</sub>. The two isomers (trans and cis) of vitamin K<sub>1</sub> are not separated. Vitamin K<sub>2</sub>-MK<sub>4</sub> and vitamin K<sub>2</sub>-MK<sub>7</sub> can be separated from vitamin K<sub>1</sub> by this method but are not yet included in the scope.

#### 4.6 Thiamine (vitamin B<sub>1</sub>)

Thiamine is a water-soluble vitamin. Thiamine can be found in a lot of food products mainly as phosphorylated forms.

Food regulations of many countries authorize the fortification of different kind of food products with molecules having a vitamin B<sub>1</sub> activity. The names of the authorized molecular forms to add in are generally linked to the type of food product. [Table 8](#) gives an example of vitamin B<sub>1</sub> compounds permitted for use for food fortification according to EU regulations, depending on the product type.

**Table 8 — Example of authorized vitamin B<sub>1</sub> compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	Thiamine hydrochloride	337,3	Regulation (EU) n° 609/2013
	Thiamine mononitrate	327,4	
Supplemented food	Thiamine hydrochloride	337,3	Regulation (EC) n° 1925/2006
	Thiamine mononitrate	327,4	
Food supplements	Thiamine hydrochloride	337,3	Directive n°2002/46/EC
	Thiamine mononitrate	327,4	
	Thiamine chloride monophosphate	380,8	
	Thiamine chloride pyrophosphate	460,8	

The commonly used unit of measurement for vitamin B<sub>1</sub> was 1 mg of thiamine hydrochloride, but nowadays it is considered to be 1 mg of thiamine base. The molecular mass of the thiamine cation is 265,3.

For expressing the different vitamin B<sub>1</sub> compounds with the different molecular masses, a correction factor is used:

- 1 mg of vitamin B<sub>1</sub> = 1 mg of thiamine base (M = 300,8);
- 1 mg of thiamine mononitrate (M = 327,4) = 0,919 mg of thiamine base (M = 300,8);
- 1 mg of thiamine hydrochloride (M = 337,3) = 0,892 mg of thiamine base (M = 300,8);
- 1 mg of thiamine ion (M = 265,3) = 1,134 mg of thiamine base (M = 300,8).

To conclude, the vitamin B<sub>1</sub> content can be calculated as:

$$\text{vitamin B}_1 (\text{mg}/100 \text{ g}) = \text{thiamine base} (\text{mg}/100 \text{ g}).$$

There are several analytical approaches for quantifying vitamin B<sub>1</sub>. Vitamin B<sub>1</sub> should be liberated as a free form. Then it can be analysed by HPLC and quantified directly by mass spectrometry, or by fluorimetry after a derivatization step.

EN 14122<sup>[18]</sup>, for example, includes an extraction of thiamine after an acid hydrolysis and an enzymatic treatment. Thiamine is separated by HPLC and quantified by fluorimetry after a pre- or post-derivatization.

ISO 21470, for example, specifies a method for the simultaneous quantitative determination of four water-soluble vitamins in infant formula and related nutritional products, including relevant forms of vitamins B<sub>1</sub>, -B<sub>2</sub>, -B<sub>3</sub> and -B<sub>6</sub> by enzymatic digestion and UHPLC-MS/MS quantification. ISO 21470 is not intended to be used on products where vitamins have not been added.

#### 4.7 Riboflavin (vitamin B<sub>2</sub>)

Riboflavin is a water-soluble vitamin. Riboflavin can be found in a lot of food products, mainly as phosphorylated forms.

Food regulations of many countries authorize the fortification of different kinds of food products with molecules having a vitamin B<sub>2</sub> activity. The names of the authorized molecular forms to add in are generally linked to the type of food product. [Table 9](#) gives an example of vitamin B<sub>2</sub> compounds permitted for use for food fortification according to EU regulations, depending on the product type.

**Table 9 — Example of authorized vitamin B<sub>2</sub> compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific dietary purpose	Riboflavin	376,4	Regulation (EU) n° 609/2013
	Riboflavin 5'phosphate sodium	478,3	
Supplemented food	Riboflavin	376,4	Regulation (EC) n° 1925/2006
	Riboflavin 5'phosphate sodium	478,3	
Food supplement products	Riboflavin	376,4	Directive n°2002/46/EC
	Riboflavin 5'phosphate sodium	478,3	

Nowadays, the commonly used unit of measurement for vitamin B<sub>2</sub> is 1 mg of riboflavin.

The vitamin B<sub>2</sub> content can be calculated as:

$$\text{vitamin B}_2 \text{ (mg/100 g)} = \text{riboflavin (mg/100 g)}.$$

It should be noted that food regulations allow the addition of riboflavin as a food additive (colouring agent) in food products with the following compounds:

- E 101 (i): Riboflavin;
- E 101 (ii): Riboflavin 5'phosphate;
- E 101 (iii): Riboflavin, from *Bacillus Subtilis*.

At the analysis step this can lead to a difference between the vitamin B<sub>2</sub> quantified value and the vitamin B<sub>2</sub> labelled value because no test is capable to identify the part of vitamin B<sub>2</sub> added for nutritional purpose from the part for colouring. In any case, the labelled vitamin B<sub>2</sub> value should not include the riboflavin added as a colouring agent.

There are several analytical approaches for quantifying vitamin B<sub>2</sub>. Riboflavin should be liberated as a free form. Then it can be analysed by HPLC and quantified directly by mass spectrometry, or by fluorimetry. There is no need of a derivatization step because riboflavin is naturally fluorescent.

EN 14152<sup>[20]</sup>, for example, includes an extraction of riboflavin after an acid hydrolysis and an enzymatic treatment. Riboflavin is separated by HPLC and quantified by fluorimetry.

ISO 21470, for example, specifies a method for the simultaneous quantitative determination of four water-soluble vitamins in infant formula and related nutritional products, including relevant forms of vitamins B<sub>1</sub>, -B<sub>2</sub>, -B<sub>3</sub> and -B<sub>6</sub> by enzymatic digestion and UHPLC-MS/MS quantification. ISO 21470 is not intended to be used on products where vitamins have not been added.

#### 4.8 Niacin (vitamin B<sub>3</sub> or vitamin PP)

Niacin is a water-soluble vitamin. Niacin can be found in a lot of food products, mainly as phosphorylated forms. There are two free forms of niacin, nicotinamide and nicotinic acid.

Food regulations of many countries authorize the fortification of different kinds of food products with molecules having a niacin activity. The names of the authorized molecular forms to add in are generally linked to the type of food product. [Table 10](#) gives an example of vitamin B<sub>3</sub> compounds permitted for use for food fortification according to EU regulations, depending on the product type.

**Table 10 — Example of authorized vitamin B<sub>3</sub> compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific dietary purpose	Nicotinamide	122,1	Regulation (EU) n° 609/2013
	Nicotinic acid	123,1	
Supplemented food	Nicotinamide	122,1	Regulation (EC) n° 1925/2006
	Nicotinic acid	123,1	
Food supplement products	Nicotinamide	122,1	Directive n°2002/46/EC
	Nicotinic acid	123,1	
	Inositol hexanicotinate	810,7	

Nowadays, the commonly used unit of measurement for niacin is 1 mg of nicotinic acid.

Certain regulations (EU, US and Codex Alimentarius) consider the tryptophan content as being niacin equivalent. This ought to be taken into account if it is mandatory: 60 mg tryptophan = 1 mg niacin equivalent.

If it is not specified in the regulation, the niacin content can be calculated as:

$$\text{niacin (mg/100 g)} = \text{nicotinic acid (mg/100 g)} + 1,008 \times \text{nicotinamide (mg/100 g)}.$$

There are several analytical approaches to quantify vitamin B<sub>3</sub>. Nicotinic acid and nicotinamide need to be liberated as a free form. Then they can be analysed by HPLC and quantified directly by mass spectrometry, or by fluorimetry after derivatization. Some methods involve a transformation step from one vitamers in the other. As such, the total niacin is directly quantified.

EN 14152<sup>[20]</sup>, for example, includes an extraction of niacin after an acid hydrolysis or an enzymatic treatment. Nicotinamide and nicotinic acid are separated by HPLC and quantified by fluorimetry after a derivatization step by UV action. Inositol hexanicotinate is not quantified by this method.

ISO 21470, for example, specifies a method for the simultaneous quantitative determination of four water-soluble vitamins in infant formula and related nutritional products, including relevant forms of vitamins B<sub>1</sub>, -B<sub>2</sub>, -B<sub>3</sub> and -B<sub>6</sub> by enzymatic digestion and UHPLC-MS/MS quantification. ISO 21470 is not intended to be used on products where vitamins have not been added.

#### 4.9 Vitamin B<sub>6</sub>

Vitamin B<sub>6</sub> is a water-soluble vitamin. Vitamin B<sub>6</sub> can be found in a lot of food products free or as phosphorylated forms. There are three vitamers of vitamin B<sub>6</sub>: pyridoxine (or pyridoxol), pyridoxal and pyridoxamine.

Food regulations of many countries authorize the fortification of different kinds of food products with molecules having a vitamin B<sub>6</sub> activity. The names of the authorized molecular forms to add in are generally linked to the type of food product. [Table 11](#) gives an example of vitamin B<sub>6</sub> compounds permitted for use for food fortification according to EU regulations, depending on the product type.

**Table 11 — Example of authorized vitamin B<sub>6</sub> compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	Pyridoxine hydrochloride	205,6	Regulation (EU) n° 609/2013
	Pyridoxine 5' phosphate	249,6	
	Pyridoxine dipalmitate	646,0	
Supplemented food	Pyridoxine hydrochloride	205,6	Regulation (EC) n° 1925/2006
	Pyridoxine 5' phosphate	249,6	
	Pyridoxine dipalmitate	646,0	
Food supplement products	Pyridoxine hydrochloride	205,6	Directive n° 2002/46/EC
	Pyridoxine 5' phosphate	249,6	
	Pyridoxal 5' phosphate	247,1	

Nowadays, the commonly used unit of measurement for vitamin B<sub>6</sub> is 1 mg of pyridoxin.

The three vitamers are considered to have the same vitamin B<sub>6</sub> activity:

- 1 mg of vitamin pyridoxine hydrochloride (M = 205,6) = 0,823 mg of vitamin B<sub>6</sub> (M = 169,1);
- 1 mg of pyridoxine 5' phosphate (M = 249,6) = 0,679 of vitamin B<sub>6</sub> (M = 169,1).

The vitamin B<sub>6</sub> content can be calculated as:

vitamin B<sub>6</sub> (mg/100 g) = pyridoxin (mg/100 g) + 1,012 pyridoxal (mg/100 g) + 1,006 pyridoxamine (mg/100 g)

There are several analytical approaches for quantifying vitamin B<sub>6</sub>. The three vitamers need to be liberated as free forms. Then they can be analysed by HPLC and quantified directly by mass spectrometry, or by fluorimetry (see for example EN 14663<sup>[21]</sup>). Some methods involve a transformation step from two vitamers into the third one. The total vitamin B<sub>6</sub> is directly quantified (see for example EN 14164<sup>[22]</sup>).

EN 14663<sup>[21]</sup>, for example, includes an extraction of vitamin B<sub>6</sub> after an acid hydrolysis or an enzymatic treatment. Pyridoxin, pyridoxamine and pyridoxal are separated by HPLC and quantified by fluorimetry. Pyridoxine dipalmitate is not quantified by this method.

EN 14164<sup>[22]</sup>, for example, includes an extraction of vitamin B<sub>6</sub> after an acid hydrolysis or an enzymatic treatment. Pyridoxamine and pyridoxal are transformed to pyridoxin. Then pyridoxin is separated by HPLC and quantified by fluorimetry. Pyridoxine dipalmitate is not quantified by this method.

ISO 21470, for example, specifies a method for the simultaneous quantitative determination of four water-soluble vitamins in infant formula and related nutritional products, including relevant forms of vitamins B<sub>1</sub>, -B<sub>2</sub>, -B<sub>3</sub> and -B<sub>6</sub> by enzymatic digestion and UHPLC-MS/MS quantification. ISO 21470 is not intended to be used on products where vitamins have not been added.

#### 4.10 Biotin (vitamin H or vitamin B<sub>8</sub>)

Biotin is a water-soluble vitamin. Biotin can be found in a lot of food products free or bound to proteins or peptides. There are two main vitamers with a biotin activity, D-biotin and D-Biocytyl. Biocytyl is rarely detected. But as intermediate in the metabolism of biotin, biocytyl occurs naturally in blood serum and can be detected in offal.

Food regulations of many countries authorize the fortification of different kinds of food products with biotin. The name of the authorized molecular form to add in is generally linked to the type of food product. [Table 12](#) gives an example of biotin permitted for use for food fortification according to EU regulations, depending on the product type.

**Table 12 — Example of authorized molecules in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	D-Biotin	244,3	Regulation (EU) n° 609/2013
Supplemented food	D-Biotin	244,3	Regulation (EC) n° 1925/2006
Food supplement products	D-Biotin	244,3	Directive n°2002/46/EC

Nowadays, the commonly used unit of measurement for biotin is 1 µg of biotin.

1 µg of D-biocytyin = 0,656 µg of biotin.

The biotin content can be calculated as:

$$\text{biotin } (\mu\text{g}/100 \text{ g}) = \text{D-Biotin } (\mu\text{g}/100 \text{ g}) + 0,656 \text{ D-Biocytyin } (\mu\text{g}/100 \text{ g})$$

D-biocytyin is rarely quantified, but should be taken in account in the calculation if it is the case.

There are several analytical approaches for quantifying biotin. Biotin should be liberated as a free form. Then it can be analysed by HPLC and quantified by fluorimetry after a derivatization step (see for example EN 15607<sup>[24]</sup>) or by UV. In this case, a preconcentration step should be added before the HPLC (see ISO 23305, for example).

In ISO 23305, for example, the sample is dispersed in a phosphate buffer and then autoclaved. Biotin is purified and concentrated with an immunoaffinity column. Then it is analysed by HPLC and UV quantified. Biocytyin and biotin are separated.

EN 15607<sup>[24]</sup>, for example, includes an enzymatic treatment. Then the extract is analysed by HPLC with a post-column derivatization. Biocytyin and biotin are separated and quantified by fluorimetry.

#### 4.11 Folates (vitamin B<sub>9</sub>)

Folic acid is the reference compound of a group of different vitamers which have a vitamin B<sub>9</sub> activity. Folic acid is a water-soluble vitamin B<sub>9</sub>. Folates can be found in a lot of food products, free or bound to chains of different numbers of glutamate units.

The main free forms are folic acid, 5-methyl-tetrahydrofolate and 5-formyl-tetrahydrofolate. But there are around ten compounds which are considered to have vitamin B<sub>9</sub> activity.

Food regulations of many countries authorize the fortification of different kinds of food products with folates. The names of the authorized molecular forms to add in are generally linked to the type of food product. [Table 13](#) gives an example of folates permitted for use for food fortification according to EU regulations depending the product type.

**Table 13 — Example of authorized molecules in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	Folic acid (pteroylmonoglutamate acid)	441,4	Regulation (EU) n° 609/2013
	Calcium L-methylfolate	459,5	
Supplemented food	Folic acid (pteroylmonoglutamate acid)	441,4	Regulation (EC) n° 1925/2006

Table 13 (continued)

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food supplements	Folic acid (pteroylmonoglutamate acid)	441,4	Directive n°2002/46/EC
	Calcium L-methylfolate	459,5	

The commonly used unit of measurement for folates used to be 1 µg of folic acid.

All folates quantified were expressed in folic acid. However, multiple regulations have now introduced Dietary Folate Equivalents (DFE) as the unit of measurement; see [Table 14](#).

Table 14 — USA unit conversion for folates

Nutrient	Former unit	Former conversion	New unit	New conversion
Folate	µg		µg DFE (Dietary Folate Equivalent)	1 µg DFE = 1 µg folates 1 µg DFE = 0,6 µg folic acid

Recently in the EU, two new regulations have introduced the same principle: one for food for special medical purpose<sup>[25]</sup> and another for infant formula<sup>[26]</sup>.

The folates (vitamin B<sub>9</sub>) content can be calculated as:

vitamin B<sub>9</sub> as DFE (µg/100 g) = 1/0,6 folic acid content + folates content (µg/100 g),

or if it is outlined by regulation, the unit of measurement of vitamin B<sub>9</sub> = µg DFE.

There are several analytical approaches for quantifying folates. Vitamin B<sub>9</sub> was historically quantified by microbiological methods. The calibration was made with folic acid. Some HPLC methods have been more recently developed. A detection and a quantification by mass spectrometry enable differentiation between the different folates. In any case, a trienzymatic digestion should be used beforehand to prepare the samples. Although the LC techniques are nowadays widely deployed for the analysis of vitamins, microbiological methods are still currently used for vitamin B<sub>9</sub> (for example EN 14131<sup>[32]</sup>). Having two completely different principles often means discrepancy in results.

In AOAC 2011.06<sup>[27]</sup> for example, the folates in a sample are extracted in a buffer (pH = 6,0) containing internal standards by treatments with protease, amylase and rat plasma conjugase (trienzyme digestion). The extract is purified and concentrated using a weak anion exchange (WAX) Solid Phase Extraction (SPE). Polyglutamate forms of folates in the sample are deconjugated to monoglutamates during the extraction and are analysed by LC-MS/MS. Folic acid, 5-methyl-tetrahydrofolate (5-CH<sub>3</sub>-THF) and 5-formyl-tetrahydrofolate (5-CHO-THF) are quantified, and total folate is estimated and expressed as folic acid. Isotopically labelled folic acid (<sup>13</sup>C-folic acid), 5-CH<sub>3</sub>-THF (<sup>13</sup>C-5-CH<sub>3</sub>-THF) and 5-CHO-THF (<sup>13</sup>C-5-CHO-THF) are used as the internal standards.

#### 4.12 Vitamin B<sub>12</sub>

Vitamin B<sub>12</sub> is a water-soluble vitamin. Vitamin B<sub>12</sub> can be found in all food from animal origin (offal, sea food, meat, fish, eggs and milk products).

Cobalamin is the reference compound of a group of different vitamers which have a vitamin B<sub>12</sub> activity. There are four vitamers of vitamin B<sub>12</sub>; cyanocobalamin, hydroxycobalamin, methylcobalamin and adenosylcobalamin.

Food regulations of many countries authorize the fortification of different kinds of food products with vitamin B<sub>12</sub>. The names of the authorized molecular forms to add in is generally linked to the type

of food product. [Table 15](#) gives an example of the vitamin B<sub>12</sub> compound permitted for use for food fortification according to European regulation depending the product type.

**Table 15 — Example of authorized vitamin B<sub>12</sub> compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	Cyanocobalamine	1 355,4	Regulation (EU) n° 609/2013
	Hydroxycobalamine	1 346,4	
Supplemented food	Cyanocobalamine	1 355,4	Regulation (EC) n° 1925/2006
	Hydroxycobalamine	1 346,4	
Food supplements	Cyanocobalamine	1 355,4	Directive n°2002/46/EC
	Hydroxycobalamine	1 346,4	

Nowadays, the commonly used unit of measurement for vitamin B<sub>12</sub> is 1 µg of cyanocobalamin.

The vitamin B<sub>12</sub> content can be calculated as:

vitamin B<sub>12</sub> (µg/100 g) = sum of the different vitamers expressed as cyanocobalamin (µg/100 g).

There are several analytical approaches for quantifying vitamin B<sub>12</sub>. Vitamin B<sub>12</sub> was historically quantified by microbiological methods. The calibration was done with cyanocobalamin. More recently, some HPLC methods have been developed. Although LC techniques are nowadays widely deployed for the analysis of vitamins, microbiological methods are still currently used for vitamin B<sub>12</sub>. Having two completely different principles often means discrepancy in results.

In ISO 20634, for example, cyanocobalamin and other cobalt-containing corrinoids are extracted from the sample using sodium acetate buffer (pH = 4,5) and the latter converted to cyanocobalamin using potassium cyanide at 105 °C. Extracts are purified and concentrated with C8 or C18 solid-phase extraction (SPE) cartridges and analysed with size-exclusion and reversed-phase chromatography. Determination of vitamin B<sub>12</sub> is made by liquid chromatography with visible detection at 550 nm.

#### 4.13 Pantothenic acid (vitamin B<sub>5</sub>)

D-Pantothenic acid is a water-soluble vitamin. D-Pantothenic acid can be found in nearly all food, with higher amounts found in offal, eggs and poultry.

Dexpanthothenol (D-pantothenol) is the alcohol analogue of D-Pantothenic acid. It is not naturally present in food but can be used for fortification.

Food regulations of many countries authorize the fortification of different kinds of food products with D-pantothenic acid. The names of the authorized molecular forms to add in are generally linked to the type of food product. [Table 16](#) gives an example of the pantothenic acid compounds permitted for use for food fortification according to EU regulations, depending on the product type.

**Table 16 — Example of authorized pantothenic acid compounds for food fortification in the EU according to products concerned**

Product type	Molecular form	Molecular mass (g/mol)	Examples of regulations
Food with a specific nutritional purpose	Calcium D-pantothenate	476,5	Regulation (EU) n° 609/2013
	Sodium D-pantothenate	241,2	
	Dexpanthothenol (D-pantothenol)	205,3	